

AMBIENT AQUATIC LIFE WATER QUALITY CRITERIA FOR
NICKEL

U.S. ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF RESEARCH AND DEVELOPMENT
ENVIRONMENTAL RESEARCH LABORATORIES
DULUTH, MINNESOTA
NARRAGANSETT, RHODE ISLAND



NOTICES

This document has been reviewed by the Criteria and Standards Division, Office of Water Regulations and Standards, U.S. Environmental Protection Agency, and approved for publication.

Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

This document is available to the public through the National Technical Information Service (NTIS), 5285 Port Royal Road, Springfield, VA 22161.

ACKNOWLEDGMENTS

Loren J. Larson
Judy L. Crane
(freshwater authors)
University of Wisconsin-Superior
Superior, Wisconsin

Jeffrey L. Hyland
Robert E. Hillman
(saltwater authors)
Battelle New England Laboratory
Duxbury, Massachusetts

Charles E. Stephan
(document coordinator)
Environmental Research Laboratory
Duluth, Minnesota

David J. Hansen
(saltwater coordinator)
Environmental Research Laboratory
Narragansett, Rhode Island

Clerical Support: Terry L. Highland
Shelley A. Heintz
Diane L. Spehar

CONTENTS

	<u>Page</u>
Foreword	iii
Acknowledgments	iv
Tables	vi
Introduction	1
Acute Toxicity to Aquatic Animals	5
Chronic Toxicity to Aquatic Animals	8
Toxicity to Aquatic Plants	10
Bioaccumulation	11
Other Data	12
Unused Data	14
Summary	16
National Criteria	17
References	54

TABLES

	<u>Page</u>
1. Acute Toxicity of Nickel to Aquatic Animals	20
2. Chronic Toxicity of Nickel To Aquatic Animals	30
3. Ranked Genus Mean Acute Values with Species Mean Acute-Chronic Ratios	33
4. Toxicity of Nickel to Aquatic Plants	38
5. Bioaccumulation of Nickel by Aquatic Organisms	40
6. Other Data on Effects of Nickel on Aquatic Organisms	42

Introduction*

Nickel is one of the most common of the metals occurring in surface waters (Forstner 1984; Hutchinson et al. 1975; Kopp and Kroner 1967; Martin and Knauer 1972; Mathis and Cummings 1973; McCabe et al. 1970; Portman 1972; Solbe 1973; Trollope and Evans 1976; Young 1982). Although nickel can exist in oxidation states of -1, 0, +1, +2, +3, and +4, under usual conditions in surface waters the divalent cation greatly predominates and is generally considered the most toxic. Alkalinity, hardness, salinity, pH, temperature, and such complexing agents as humic acids influence the oxidation state, toxicity, and availability of the nickel in aquatic ecosystems.

Natural sources of the nickel in surface waters include weathering of rocks, inflow of particulate matter, and precipitation. Anthropogenic sources of nickel include the burning of coal and other fossil fuels and discharges from such industries as electroplating and smelting. Although fly ash can contain as much as 960 $\mu\text{g/g}$ (Swaine 1980), lake restoration projects have experimented with the use of fly ash to remove nutrients.

Mechanisms of nickel toxicity are varied and complex (Mushak 1980), and, as with other heavy metals, significant effects occur at cell membranes and membranous tissues, such as gills. In fish, hematological effects such as hyperglycemia, lymphopenia, and erythrocytosis have been reported in association with nickel intoxication (Agrawal et al. 1979; Chaudhry 1984; Chaudhry and Nath 1985; Chaudhry and Nath 1985; Gill and Pant 1981).

* An understanding of the "Guidelines for Deriving Numerical National Water Quality Criteria for the Protection of Aquatic Organisms and Their Uses" (Stephan et al. 1985), hereafter referred to as the Guidelines, and the response to public comment (U.S. EPA 1985a) is necessary in order to understand the following text, tables, and calculations.

Because of the variety of forms of nickel (Callahan et al. 1979; Nriagu 1980) and lack of definitive information about their relative toxicities, no available analytical measurement is known to be ideal for expressing aquatic life criteria for nickel. Previous aquatic life criteria for nickel (U.S. EPA 1980) were expressed in terms of total recoverable nickel (U.S. EPA 1983a), but this measurement is probably too rigorous in some situations. Acid-soluble nickel (operationally defined as the nickel that passes through a 0.45 μm membrane filter after the sample is acidified to pH = 1.5 to 2.0 with nitric acid) is probably the best measurement at present for the following reasons:

1. This measurement is compatible with nearly all available data concerning toxicity of nickel to, and bioaccumulation of nickel by, aquatic organisms. No test results were rejected just because it was likely that they would have been substantially different if they had been reported in terms of acid-soluble nickel. For example, results reported in terms of dissolved nickel would not have been used if the concentration of precipitated nickel had been substantial.
2. On samples of ambient water, measurement of acid-soluble nickel will probably measure all forms of nickel that are toxic to aquatic life or can be readily converted to toxic forms under natural conditions. In addition, this measurement probably will not measure several forms, such as nickel that is occluded in minerals, clays, and sand or is strongly sorbed to particulate matter, that are not toxic and are not likely to become toxic under natural conditions. Although this measurement (and many others) will measure soluble complexed forms of nickel, such as the EDTA complex of nickel, that probably have low toxicities

to aquatic life, concentrations of these forms probably are negligible in most ambient water.

3. Although water quality criteria apply to ambient water, the measurement used to express criteria is likely to be used to measure nickel in aqueous effluents. Measurement of acid-soluble nickel probably will be applicable to effluents because it will measure precipitates, such as carbonate and hydroxide precipitates of nickel, that might exist in an effluent and dissolve when the effluent is diluted with receiving water. If desired, dilution of effluent with receiving water before measurement of acid-soluble nickel might be used to determine whether the receiving water can decrease the concentration of acid-soluble nickel because of sorption.
4. The acid-soluble measurement is probably useful for most metals, thus minimizing the number of samples and procedures that are necessary.
5. The acid-soluble measurement does not require filtration at the time of collection, as does the dissolved measurement.
6. The only treatment required at the time of collection is preservation by acidification to pH = 1.5 to 2.0, similar to that required for the total recoverable measurement.
7. Durations of 10 minutes to 24 hours between acidification and filtration of most samples of ambient water probably will not affect the result substantially.
8. The carbonate system has a much higher buffer capacity from pH = 1.5 to 2.0 than it does from pH = 4 to 9 (Weber and Stumm 1963).
9. Differences in pH within the range of 1.5 to 2.0 probably will not affect the result substantially.

10. The acid-soluble measurement does not require a digestion step, as does the total recoverable measurement.
 11. After acidification and filtration of the sample to isolate the acid-soluble nickel, the analysis can be performed using either atomic absorption spectrophotometric or ICP-atomic emission spectrometric analysis (U.S. EPA 1983a), as with the total recoverable measurement.
- Thus, expressing aquatic life criteria for nickel in terms of the acid-soluble measurement has both toxicological and practical advantages. On the other hand, because no measurement is known to be ideal for expressing aquatic life criteria for nickel or for measuring nickel in ambient water or aqueous effluents, measurement of both acid-soluble nickel and total recoverable nickel in ambient water or effluent or both might be useful. For example, there might be cause for concern if total recoverable nickel is much above an applicable limit, even though acid-soluble nickel is below the limit.
- Unless otherwise noted, all concentrations reported herein are expected to be essentially equivalent to acid-soluble nickel concentrations. All concentrations are expressed as nickel, not as the chemical tested. The criteria presented herein supersede previous national aquatic life water quality criteria for nickel (U.S. EPA 1976, 1980) because these new criteria were derived using improved procedures and additional information. Whenever adequately justified, a national criterion may be replaced by a site-specific criterion (U.S. EPA 1983b), which may include not only site-specific criterion concentrations (U.S. EPA 1983c), but also site-specific durations of averaging periods and site-specific frequencies of allowed excursions (U.S. EPA 1985b). The latest comprehensive literature search for information for this document was conducted in July, 1986; some more recent information might have been included.

Acute Toxicity to Aquatic Animals

Lind et al. (Manuscript) conducted studies on the effects of both hardness and TOC on the acute toxicity of nickel to both Daphnia pulicaria and the fathead minnow (Table 6). With both species, hardness was the only significantly correlated parameter. Nebeker et al. (1985) reported that rainbow trout were more sensitive when 12-months old than when 3-months old. Rehwoldt et al. (1973) observed that embryos were more sensitive than adult snails.

Many factors might affect the results of tests of the toxicity of nickel to aquatic organisms (Sprague 1985), but water quality criteria can quantitatively take into account such factors only if enough data are available to show that the factor similarly affects the results of tests with a variety of species. Hardness is often thought of as having a major effect on the toxicity of nickel in fresh water, although the observed effect is probably due to one or more of a number of usually interrelated ions, such as hydroxide, carbonate, calcium, and magnesium. Hardness (expressed as mg CaCO₃/L) is used here as a surrogate for the ions that affect the results of toxicity tests on nickel. An analysis of covariance (Dixon and Brown 1979; Neter and Wasserman 1974) was performed using the natural logarithm of the acute value as the dependent variable, species as the treatment or grouping variable, and the natural logarithm of hardness as the covariate or independent variable. This analysis of covariance model was fit to the data in Table 1 for the four species for which acute values are available over a range of hardness such that the highest hardness is at least three times the lowest and the highest is also at least 100 mg/L higher than the lowest. The four slopes are between 0.69 and 1.19 (see end of Table 1) and are close to the slope

of 1.0 that is expected on the basis that nickel, calcium, magnesium, and carbonate all have a charge of two. An F-test showed that, under the assumption of equality of slopes, the probability of obtaining four slopes as dissimilar as these is $P = 0.26$. This was interpreted as indicating that it is not unreasonable to assume that the slopes for these four species are the same.

Where possible, the pooled slope of 0.8460 was used to adjust the acute values in Table 1 to hardness = 50 mg/L. Species Mean Acute Values were calculated as geometric means of the adjusted acute values. Genus Mean Acute Values at hardness = 50 mg/L were then calculated as geometric means of the available freshwater Species Mean Acute Values (Table 3). Of the eighteen genera for which freshwater acute values are available, the most sensitive genus, Daphnia, was 29 times more sensitive than the most resistant, Fundulus. The freshwater Final Acute Value for nickel at hardness = 50 mg/L was calculated to be 1,578 $\mu\text{g}/\text{L}$ using the procedure described in the Guidelines and the Genus Mean Acute Values in Table 3. Thus, the freshwater Criterion Maximum Concentration (in $\mu\text{g}/\text{L}$) =
$$(0.8460[\ln(\text{hardness})]+3.3612)$$

The acute toxicity of nickel to saltwater organisms has been determined with 18 species of invertebrates and 4 species of fish (Table 1). The LC50s and EC50s for invertebrates range from 151.7 $\mu\text{g}/\text{L}$ for juveniles of the mysid, Heteromysis formosa (Gentile et al. 1982) to 1,100,000 $\mu\text{g}/\text{L}$ for late juvenile to adult stages of the clam, Macoma balthica (Bryant et al. 1985). Fish are not as sensitive or as resistant to nickel. The 96-hr LC50s range from 7,958 $\mu\text{g}/\text{L}$ for larval stages of the Atlantic silverside, Menidia menidia (Cardin 1985) to 350,000 $\mu\text{g}/\text{L}$ for adult stages of the mummichog, Fundulus heteroclitus (Eisler and Hennekey 1977).

Although data are limited, relationships might exist between both salinity and temperature and the toxicity of nickel to some saltwater species. For example, the LC50 for the mummichog is 55,000 µg/L at a salinity of 6.9 g/kg, and 175,000 µg/L at a salinity of 21.6 g/kg (Dorfman 1977). In a series of tests with the amphipod, Corophium volutator (Bryant et al. 1985), the LC50 increased with salinity at 5°C, 10°C, and 15°C. At salinities of 5, 10, and 15 g/kg, temperature did not seem to affect the LC50, but at salinities of 25 and 35 g/kg, the LC50 decreased as temperature increased. Bryant et al. (1985) found similar effects of salinity and temperature on nickel toxicity with the clam, Macoma balthica (Table 6). Regressions of toxicity on salinity for the above data show strong correlations. However, analysis of covariance reveals that the slopes for the individual species are too dissimilar ($P < 0.05$) to justify expressing nickel toxicity as a function of salinity.

Of the twenty saltwater genera for which acute values are available, the most sensitive genus, Heteromysis, was over 2,000 times more sensitive than the most resistant, Mya (Table 3). Acute values are available for more than one species in each of three genera, and the range of Species Mean Acute Values within each genus is less than a factor of 4.8. Genus Mean Acute Values for the four most sensitive genera, Heteromysis, Mercenaria, Mysidopsis, and Crassostrea, were within a factor of 7.8 even though the acute tests were conducted with juveniles of the crustaceans and with embryos of the bivalves. The saltwater Final Acute Value was calculated to be 149.2 µg/L, which is very close to the acute value for the most sensitive tested saltwater species.

Chronic Toxicity to Aquatic Animals

Data are available on the freshwater chronic toxicity of nickel to a cladoceran, a caddisfly, and two species of fish (Table 2). Nebeker et al. (1985) conducted two early life-stage tests beginning with rainbow trout embryos 4 hours after fertilization and one early life-stage test beginning with trout embryos 25 days after fertilization. In the first test, weight was significantly reduced by all tested concentrations including the lowest of 35 µg/L. In the second test, weight was significantly reduced by 62 and 431 µg/L, but not by 35, 134, and 238 µg/L, whereas survival was reduced only at nickel concentrations of 134 µg/L and higher. In the third test, weight was significantly reduced at 431 µg/L and higher, but the reduction in survival was significant only at 1,680 µg/L and higher.

Lazareva (1985) conducted a life-cycle test over successive generations with Daphnia magna and observed little change in sensitivity. Although survival time was the most sensitive parameter in one test, growth was consistently affected at a concentration of 10 µg/L. Lazareva predicted that 5 µg/L would affect the productivity of populations of Daphnia magna.

The influence of hardness on chronic toxicity of nickel was investigated by Chapman et al. (Manuscript). In life-cycle tests with Daphnia magna, they observed an increase in chronic value with increased hardness. Least squares regression of ln[chronic value] on ln[hardness] produced a slope of 2.3007 with wide confidence limits (Table 2). A similar regression with data for the fathead minnow produced a slope of 0.5706, but confidence limits could not be calculated because only two points were available for use in the regression. An F-test showed that, under the

assumption of equality of slopes, the probability of obtaining two slopes as dissimilar as these is $P = 0.19$. This was interpreted as indicating that it is not unreasonable to assume that the two slopes are the same. The pooled slope is 1.3418 with 95% confidence limits of -1.3922 and 4.0760. The confidence limits on the pooled acute slope are well within the confidence limits on the pooled chronic slope.

The mysid, Mysidopsis bahia, is the only saltwater species with which an acceptable chronic test has been conducted on nickel (Table 2). Chronic exposure to nickel reduced survival and number of young at 141 $\mu\text{g/L}$ and above but not at 61 $\mu\text{g/L}$ and lower (Lussier et al. 1985). Thus the chronic value for nickel with this species is 92.74 $\mu\text{g/L}$ and the acute-chronic ratio is 5.478.

The three available species mean acute-chronic ratios range from 5.478 to 35.58 and were all determined with species that are acutely sensitive to nickel (Table 3). The Final Acute-Chronic Ratio of 17.99 was calculated as the geometric mean of the three ratios. Division of the freshwater Final Acute Value by the Final Acute-Chronic Ratio results in a freshwater Final Chronic Value of 87.72 $\mu\text{g/L}$ at hardness = 50 mg/L. Some data (Tables 2 and 6) concerning the chronic toxicity of nickel to rainbow trout indicate that embryos and larvae of this species might be affected at this concentration, whereas other data (Table 2) indicate that embryos and larvae of the species might not be adversely affected. Use of an acute-chronic ratio that is independent of hardness is equivalent to assuming that the chronic slope is equal to the acute slope. Thus the freshwater Final Chronic Value (in $\mu\text{g/L}$) = $e^{(0.8460[\ln(\text{hardness})]+1.645)}$. This value might not protect Daphnia magna in soft water.

Division of the saltwater Final Acute Value by 17.99 results in a saltwater Final Chronic Value of 8.293 µg/L, which is about a factor of eleven lower than the only chronic value that has been determined with a saltwater species. Three of the four acutely most sensitive saltwater species are in the same family as the species with which the saltwater acute-chronic ratio was determined. In addition, two other sensitive species are bivalve molluscs for which the acute values were obtained from tests on embryos and larvae.

Toxicity to Aquatic Plants

Data on the toxicity of nickel to aquatic plants are found in Table 4. Nickel concentrations resulting in a 40-60% reduction in growth of freshwater algae range from 50 µg/L for the green alga, Scenedesmus acuminatus, to 5,000 µg/L for the green algae, Ankistrodesmus falcatus and Chlorococcum sp. Wang and Wood (1984) indicate that toxicity of nickel to plants is pH dependent. Although lack of hardness values makes comparisons difficult, general comparison of data in Table 4 with chronic toxicity data in Table 2 suggests that nickel concentrations high enough to produce chronic effects on freshwater animals will also have deteriorative effects on freshwater algal populations.

Patrick et al. (1975) found that nickel decreased diatom diversity and caused a shift to green and blue-green algae. In their field study, Spencer and Greene (1981) also found an increase in blue-green algae. Using EDTA to manipulate Ni⁺² concentrations, Spencer and Nichols (1983) reported algal growth to be inversely related to free divalent nickel and independent of total nickel concentrations.

Watras et al. (1985) reported a BCF with Daphnia magna of 11.6. Their study indicated that uptake of nickel directly from the water was much greater than uptake from food. They also suggested that little biomagnification occurs within the association of the cladoceran and algae. Jennett et al. (1982) examined physical and biological variables affecting uptake by algae. Although their study did not demonstrate that steady-state was attained, Taylor and Crowder (1983,1984) studied differential uptake of nickel by various portions of an emergent aquatic plant, the cattail. A field study with measured nickel concentrations in a stream produced average BCF of 803 for wild rainbow trout (Salmo gairdneri) (Dallinger and Kautzky 1985).

Data on bioaccumulation of nickel by saltwater organisms are available for two species of algae and two species of bivalves (Table 5). BCFs for algae collected from the field are 675 for the rockweed, Fucus vesiculosus, and 458.3 for Ascophyllum nodosum (Foster 1976). BCFs for bivalves exposed for 9 days in the laboratory were 472.7 and 328.6 for the blue mussel and 458.1 and 261.8 for the Eastern oyster (Zaroogian and Johnson 1984).

No U.S. FDA action level or other maximum acceptable concentration in tissue is available for nickel, and, therefore, no Final Residue Value can be calculated.

Other Data

Data in Table 6 suggest a high toxicity to nickel in the single-celled organisms. Bringmann and Kuhn (1959a,b;1977a;1979;1980a,b;1981) reported that concentrations of 2.5 to 1,500 µg/L resulted in incipient inhibition of algae, bacteria, and protozoans. Babich and Stotzky (1983) observed delayed effects after a 24-hr exposure.

Willford (1966) reported 48-hr LC50s for six fishes tested in the same water. Although the fish differed in size, neither this nor taxonomic differences produced a clear trend in relative toxicity. Blaylock and Frank (1979) observed LC50s for carp larva at 3 and 10.5 days to be 8,460 and 750 $\mu\text{g}/\text{L}$, respectively. Birge and coworkers obtained 28-day EC50s of 50, 60, and 90 $\mu\text{g}/\text{L}$ with embryos and larvae of rainbow trout and a 7-day EC50 of 50 $\mu\text{g}/\text{L}$ with embryos and larvae of the narrow-mouthed toad.

Shaw and Brown (1971) studied the effect of nickel on laboratory fertilization of rainbow trout eggs. They did not find a statistically significant effect at 1000 $\mu\text{g}/\text{L}$ (hardness = 260 to 280 mg/L) and noted a stimulation in development after fertilization compared to controls.

Whitley and Sikora (1970) and Brkovic-Popovic and Popovic (1977b) studied effects on respiration in tubificid worms. Influence of nickel on thermal resistance of salmonids was examined by Becker and Wolford (1980). The effect of complexing agents on toxicity of nickel to carp was studied by Muramoto (1983). Smith-Sonneborn et al. (1983) studied the toxicity of nickel dust particles ingested by Paramecium. Anderson (1973) and Anderson and Weber (1975) derived an expression relating body size to sensitivity of the guppy.

In a field study, Havas and Hutchinson (1982) worked with acidified and control ponds and suggested that the lowered pH increased the concentrations of heavy metals such as nickel and stressed resident aquatic invertebrates. Keller and Pitblado (1984) and Yan et al. (1985) compared ambient nickel concentrations to aquatic community dynamics.

Available data that were not used directly in the derivation of saltwater criterion for nickel (Table 6) do not indicate a need to lower the criterion. In addition to affecting survival of saltwater animals,

nickel affects growth, development, reproduction, and biochemical responses. A 19% reduction in growth of juvenile Pacific oysters, Crassostrea gigas, exposed to 10 $\mu\text{g}/\text{L}$ for 14 days at a salinity of 34 g/kg was reported by Watling (1983). The ecological significance of this reduction is unknown, but after 14 days in clean water size was similar to that of the controls. Petrich and Reish (1979) found that 100 to 500 $\mu\text{g}/\text{L}$ suppressed reproduction of a polychaete, Ctenodrilus serratus. Zaroogian et al. (1982) showed a significant reduction in ATP activity in the adductor muscle of the blue mussel, but not the Eastern oyster, after a 10-week exposure to 10 $\mu\text{g}/\text{L}$. Abnormal development of embryos of the sea urchins, Arbacia punctulata and Lytocrinus pictus, occurred at several concentrations of nickel (Timourian and Watchmaker 1972; Waterman 1937), and concentrations as low as 58.69 $\mu\text{g}/\text{L}$ depressed sperm motility in gametes of the purple urchin, Strongylocentrotus purpuratus (Timourian and Watchmaker 1977).

Unused Data

Some data on the effects of nickel on aquatic organisms were not used because the studies were conducted with species that are not resident in North America (e.g., Ahsanullah 1982; Ballester and Castellvi 1979; Baudouin and Scoppa 1974; Kanai and Wakabayashi 1984; Khangarot et al. 1982; McFeters et al. 1983; Saxena and Parashari 1983; Srivastava et al. 1985; Van Hoof and Nauwelaers 1984; Verma et al. 1981; Wilson 1983). Results (e.g., Kiss et al. 1984) of tests conducted with brine shrimp, Artemia sp., were not used because these species are from a unique saltwater environment. Data were also not used if nickel was a component of a mixture (e.g., Alman and Bager 1984; Anderson 1983; Besser 1985; Cowgill et al. 1986; Doudoroff 1956; Doudoroff et al. 1966; Eisler 1977b; Hutchinson and

Sprague 1983,1986; Lopez-Avila et al. 1985; Markarian et al. 1980; Muska 1978; Muska and Weber 1977a,b; Phelps et al. 1981; Suloway et al. 1983; Stratton and Corke 1979b; Vymazal 1984; Weinstein and Anderson 1978; Wong and Beaver 1980; Wong et al. 1978,1982), an effluent (e.g., Abbe 1982; Blaise and Couture 1984; Cherry et al. 1979; Jay and Muncy 1979; Lewis 1986) or sediments (e.g., Malueg et al. 1984; Seeleye et al. 1982).

Babich and Stotzky (1985), Biddinger and Gloss (1984), Birge and Black (1980), Chapman et al. (1968), Doudoroff and Katz (1953), Eisler (1981), Jenkins (1980), Kaiser (1980), LeBlanc (1984), McKim (1977), Phillips and Russo (1978), Rai et al. (1981), Thompson et al. (1972), and U.S. EPA (1975) only contain data that have been published elsewhere. Christensen et al. (1985) reported computer simulated data only.

Data were not used if the organisms were exposed to nickel in food (e.g., Cowgill et al. 1985; Mansouri-Aliabadi and Sharp 1985; Windom et al. 1982). Results were not used if the test procedures were not adequately described (e.g., Bean and Harris 1977; Braginskiy and Shcherban 1978; Brown 1968; Jones 1939; Petukhov and Ninonenko 1982; See et al. 1974,1975; Shcherban 1977; Sirover and Loeb 1976; Soeder and Engelmann 1984; Wang et al. 1984). The 96-hr values reported by Buikema et al. (1974a,b) were subject to error because of possible reproductive interactions (Buikema et al. 1977). Michnowicz and Weak (1984) conducted tests at too low a pH. Babich et al. (1986) only exposed cell cultures.

Results of some laboratory tests were not used because the tests were conducted in distilled or deionized water without addition of appropriate salts (e.g., Jones 1935; MacDonald et al. 1980; Shaw and Grushkin 1957) or were conducted in chlorinated or "tap" water (e.g., Grande and Andersen 1983; Janauer 1985). Dilution waters in studies by Mann and Fyfe (1984)

and Stratton and Corke (1979a) contained excessive amounts of EDTA. Stokes (1975) and Whitton and Shehata (1982) used algae from waters containing high concentrations of nickel. The data of Gerhards and Weller (1977) on accumulation of nickel by algae were not used because the test concentrations of nickel adversely affected the growth of the algae. Dugan (1975) reported results in uptake studies only as counts of radio-labeled nickel.

Bringmann and Kuhn (1982) cultured Daphnia magna in one water and conducted tests in another. Tests conducted with too few test organisms (e.g., Applegate et al. 1957; Tarzwell and Henderson 1960) were not used.

Reports of the concentrations of nickel in wild aquatic organisms (e.g., Abo-Rady 1979; Amemiya and Nakayama 1984; Bailey and Stokes 1985; Bosserman 1985; Bradley and Morris 1986; Brezina and Arnold 1977; Bryan et al. 1983; Chapman 1985; Chassard-Bouchard and Balvay 1978; Dunstan et al. 1980; Eisler et al. 1978; Gordon et al. 1980; Guilizzoni 1980; Hall et al. 1978; Heit and Klasek 1985; Jenkins 1980; Kawamata et al. 1983; La Touche and Mix 1982; McDermott et al. 1976; McHardy and George 1985; Martin 1979; Mathis and Cummings 1973; Mears and Eisler 1977; O'Conner 1976; Ozimek 1985; Parsons et al. 1972; Pennington et al. 1982; Pulich 1980; Reynolds 1979; Stokes et al. 1985; Tong et al. 1974; Trollope and Evans 1976; Uthe and Bligh 1971; Van Coille and Rousseau 1974; Wachs 1982; Wehr and Whitton 1983; Wren et al. 1983) were not used to calculate bioaccumulation factors due to the absence or insufficient number of measurements of nickel in water.

Summary

Acute values with twenty-one freshwater species in 18 genera range from 1,101 $\mu\text{g/L}$ for a cladoceran to 43,240 $\mu\text{g/L}$ for a fish. Fishes and

invertebrates are both spread throughout the range of sensitivity. Acute values with four species are significantly correlated with hardness.

Data are available concerning the chronic toxicity of nickel to two invertebrates and two fishes in fresh water. Data available for two species indicate that chronic toxicity decreases as hardness increases. The measured chronic values ranged from 14.77 $\mu\text{g}/\text{L}$ with Daphnia magna in soft water to 526.7 $\mu\text{g}/\text{L}$ with the fathead minnow in hard water. Five acute-chronic ratios are available for two species in soft and hard water and range from 14 to 122.

Nickel appears to be quite toxic to freshwater algae, with concentrations as low as 50 $\mu\text{g}/\text{L}$ producing significant inhibition. Bioconcentration factors for nickel range from 0.8 for fish muscle to 193 for a cladoceran.

Acute values for twenty-three saltwater species in twenty genera range from 151.7 $\mu\text{g}/\text{L}$ with juveniles of a mysid to 1,100,000 $\mu\text{g}/\text{L}$ with juveniles and adults of a clam. The acute values for the four species of fish range from 7,598 to 350,000 $\mu\text{g}/\text{L}$. The acute toxicity of nickel appears to be related to salinity, but the form of the relationship appears to be species-dependent.

Mysidopsis bahia is the only saltwater species with which an acceptable chronic test has been conducted on nickel. Chronic exposure to 141 $\mu\text{g}/\text{L}$ and greater resulted in reduced survival and reproduction. The measured acute-chronic ratio was 5.478.

Bioconcentration factors in salt water range from 261.8 with a oyster to 675 with a brown alga.

National Criteria

The procedures described in the "Guidelines for Deriving Numerical National Water Quality Criteria for the Protection of Aquatic Organisms

"and Their Uses" indicate that, except possibly where a locally important species is very sensitive, freshwater aquatic organisms and their uses should not be affected unacceptably if the four-day average concentration (in $\mu\text{g}/\text{L}$) of nickel does not exceed the numerical value given by $(0.8460[\ln(\text{hardness})]+1.1645)$ more than once every three years on the average and if the one-hour average concentration (in $\mu\text{g}/\text{L}$) does not exceed the numerical value given by $(0.8460[\ln(\text{hardness})]+3.3612)$ more than once every three years on the average. For example, at hardnesses of 50, 100, and 200 mg/L as CaCO_3 the four-day average concentrations of nickel are 88, 160, and 280 $\mu\text{g}/\text{L}$, respectively, and the one-hour average concentrations are 790, 1400, and 2500 $\mu\text{g}/\text{L}$.

The procedures described in the "Guidelines for Deriving Numerical National Water Quality Criteria for the Protection of Aquatic Organisms and Their Uses" indicate that, except possibly where a locally important species is very sensitive, saltwater aquatic organisms and their uses should not be affected unacceptably if the four-day average concentration of nickel does not exceed 8.3 $\mu\text{g}/\text{L}$ more than once every three years on the average and if the one-hour average concentration does not exceed 75 $\mu\text{g}/\text{L}$ more than once every three years on the average.

"Acid-soluble" is probably the best measurement at present for expressing criteria for metals and the criteria for nickel were developed on this basis. However, at this time, no EPA approved method for such a measurement is available to implement criteria for metals through the regulatory programs of the Agency and the States. The Agency is considering development and approval of a method for a measurement such as "acid-soluble." Until one is approved, however, EPA recommends applying criteria for metals using the total recoverable method. This has two impacts: (1) certain

species of some metals cannot be measured because the total recoverable method cannot distinguish between individual oxidation states, and (2) in some cases these criteria might be overly protective when based on the total recoverable method.

Three years is the Agency's best scientific judgment of the average amount of time aquatic ecosystems should be provided between excursions (U.S. EPA 1985b). The resiliencies of ecosystems and their abilities to recover differ greatly, however, and site-specific allowed excursion frequencies may be established if adequate justification is provided.

Use of criteria for developing water quality-based permit limits and for designing waste treatment facilities requires selection of an appropriate wasteload allocation model. Dynamic models are preferred for the application of these criteria (U.S. EPA 1985b). Limited data or other considerations might make their use impractical, in which case one must rely on a steady-state model (U.S. EPA 1986).

Table 1. Acute Toxicity of Nickel to Aquatic Animals

<u>Species</u>	<u>Method*</u>	<u>Chemical</u>	<u>Hardness (mg/L as CaCO_3)</u>	<u>LC50 or EC50 ($\mu\text{g/L}$)**</u>	<u>Adjusted LC50 or EC50 ($\mu\text{g/L}$)***</u>	<u>Species Mean Acute Value ($\mu\text{g/L}$)****</u>	<u>Reference</u>
<u>FRESHWATER SPECIES</u>							
Worm, <u>Nais</u> sp.	S, M	-	50	14,100	14,100	14,100	Rehwoldt et al. 1975
Snail (embryo), <u>Ammicola</u> sp.	S, M	-	50	11,400	11,400	-	Rehwoldt et al. 1975
Snail (adult), <u>Ammicola</u> sp.	S, M	-	50	14,300	14,300	12,770	Rehwoldt et al. 1975
Cladoceran, <u>Daphnia magna</u>	S, U	Nickel chloride	-	<317	-	-	Anderson 1948
Cladoceran, <u>Daphnia magna</u>	S, U	Nickel chloride	45.3	510	554.4	-	Blesinger and Christensen 1972
Cladoceran, <u>Daphnia magna</u>	S, M	Nickel nitrate	51.1	915	898.3	-	Call et al. 1983
Cladoceran, <u>Daphnia magna</u>	S, M	Nickel chloride	51	1,800	1,770	-	Chapman et al. Manuscr lpt
Cladoceran, <u>Daphnia magna</u>	S, M	Nickel chloride	100	2,360	1,313	-	Chapman et al. Manuscr lpt
Cladoceran, <u>Daphnia magna</u>	S, M	Nickel chloride	104	1,920	1,033	-	Chapman et al. Manuscr lpt
Cladoceran, <u>Daphnia magna</u>	S, M	Nickel chloride	206	4,970	1,500	1,102	Chapman et al. Manuscr lpt
Cladoceran, <u>Daphnia pulicaria</u>	S, M	Nickel sulfate	48	2,182	2,259	-	Lind et al. Manuscr lpt
Cladoceran, <u>Daphnia pulicaria</u>	S, M	Nickel sulfate	48	1,813	1,877	-	Lind et al. Manuscr lpt
Cladoceran, <u>Daphnia pulicaria</u>	S, M	Nickel sulfate	44	1,836	2,046	-	Lind et al. Manuscr lpt

Table 1. (Continued)

Species	Method [#]	Chemical	Hardness (mg/L as CaCO ₃)	LC50 or EC50 ($\mu\text{g/L}$) [†]	Adjusted LC50 or EC50 ($\mu\text{g/L}$) [‡]	Species Mean Acute Value ($\mu\text{g/L}$) [§]	Reference
<u>Cladoceran, <i>Daphnia pulex</i></u>	S, M	Nickel sulfate	47	1,901	2,003	2,042	Lind et al. Manuscript
<u>Amphipod, <i>Gammarus</i> sp.</u>	S, M	-	50	13,000	13,000	13,000	Rehwoldt et al. 1973
<u>Mayfly, <i>Ephemera subvaria</i></u>	S, U	Nickel sulfate	42	4,000	4,636	4,636	Warnick and Bell 1969
<u>Damsel fly, Unidentified sp.</u>	S, M	-	50	21,200	21,200	21,200	Rehwoldt et al. 1973
<u>Stonefly, <i>Acroneuria lycorias</i></u>	S, U	Nickel sulfate	40	35,500	40,460	40,460	Warnick and Bell 1969
<u>Caddisfly, Unidentified sp.</u>	S, M	-	50	30,200	30,200	30,200	Rehwoldt et al. 1973
<u>American eel, <i>Anquilla rostrata</i></u>	S, M	Nickel nitrate	53	13,000 [†]	12,370	-	Rehwoldt et al. 1971
<u>American eel, <i>Anquilla rostrata</i></u>	S, M	-	55	13,000	11,990	12,180	Rehwoldt et al. 1972
<u>Rainbow trout (2 mos), <i>Salmo gairdneri</i></u>	F, M	Nickel nitrate	-	35,500	-	-	Hale 1977
<u>Rainbow trout (juvenile), <i>Salmo gairdneri</i></u>	F, M	Nickel sulfate	-	20,100 [†]	-	-	Anderson 1981
<u>Rainbow trout (juvenile), <i>Salmo gairdneri</i></u>	F, M	Nickel sulfate	-	12,700 [†]	-	-	Anderson 1981
<u>Rainbow trout (juvenile), <i>Salmo gairdneri</i></u>	F, M	Nickel sulfate	-	28,000 [†]	-	-	Anderson 1981
<u>Rainbow trout (juvenile), <i>Salmo gairdneri</i></u>	F, M	Nickel sulfate	-	30,900 [†]	-	-	Anderson 1981
<u>Rainbow trout (juvenile), <i>Salmo gairdneri</i></u>	F, M	Nickel sulfate	-	16,900 [†]	-	-	Anderson 1981

Table 1. (continued)

<u>Species</u>	<u>Method*</u>	<u>Chemical</u>	<u>Hardness (mg/L as CaCO₃)</u>	<u>LC50 or EC50 ($\mu\text{g/L}$)^{a,b}</u>	<u>Adjusted LC50 or EC50 ($\mu\text{g/L}$)^{a,b}</u>	<u>Species Mean Acute Value ($\mu\text{g/L}$)^{a,b}</u>	<u>Reference</u>
Rainbow trout (juvenile), <u>Salmo gairdneri</u>	F, M	Nickel sulfate	-	15,900 ^f	-	-	Anderson 1981
Rainbow trout (juvenile), <u>Salmo gairdneri</u>	F, M	Nickel sulfate	-	11,300 ^f	-	-	Anderson 1981
Rainbow trout (juvenile), <u>Salmo gairdneri</u>	F, M	Nickel sulfate	-	11,100 ^f	-	-	Anderson 1981
Rainbow trout (3 mos), <u>Salmo gairdneri</u>	F, M	Nickel chloride	27- 39	10,000 14,210	10,000 14,210	-	Nebeker et al. 1985
Rainbow trout (3 mos), <u>Salmo gairdneri</u>	F, M	Nickel chloride	27- 39	10,900 15,490	10,900 15,490	-	Nebeker et al. 1985
Rainbow trout (12 mos), <u>Salmo gairdneri</u>	F, M	Nickel chloride	27- 39	8,900 12,650	8,900 12,650	-	Nebeker et al. 1985
Rainbow trout (12 mos), <u>Salmo gairdneri</u>	F, M	Nickel chloride	27- 39	8,100 11,510	8,100 11,510	13,380	Nebeker et al. 1985
Goldfish (1-2 g), <u>Carassius auratus</u>	S, U	Nickel chloride	20	9,820 21,320	9,820 21,320	21,320	Pickerling and Henderson 1966
Common carp (<20 cm), <u>Cyprinus carpio</u>	S, M	Nickel nitrate	53	10,600 ^f	10,090	-	Rehwoldt et al. 1971
Common carp, <u>Cyprinus carpio</u>	S, M	-	55	10,400	9,594	9,839	Rehwoldt et al. 1972
Fathead minnow (1-2 g), <u>Pimephales promelas</u>	S, U	Nickel chloride	20	5,180 11,250	5,180 11,250	-	Pickerling and Henderson 1966
Fathead minnow (1-2 g), <u>Pimephales promelas</u>	S, U	Nickel chloride	20	4,580 9,943	4,580 9,943	-	Pickerling and Henderson 1966
Fathead minnow (1-2 g), <u>Pimephales promelas</u>	S, U	Nickel chloride	360	42,400 7,981	42,400 7,981	-	Pickerling and Henderson 1966
Fathead minnow (1-2 g), <u>Pimephales promelas</u>	S, U	Nickel chloride	360	44,500 8,376	44,500 8,376	-	Pickerling and Henderson 1966

Table 1. (continued)

<u>Species</u>	<u>Method*</u>	<u>Chemical</u>	<u>Hardness (mg/L as CaCO₃)</u>	<u>LC50 or EC50 (μg/L)***</u>	<u>Adjusted LC50 or EC50 (μg/L)***</u>	<u>Species Mean Acute Value (μg/L)****</u>	<u>Reference</u>
Fathead minnow (immature), <i>Pimephales promelas</i>	S, U	Nickel chloride	210	27,000	8,019	-	Pickerling 1974
Fathead minnow (immature), <i>Pimephales promelas</i>	S, M	Nickel chloride	210	32,200	9,563	-	Pickerling 1974
Fathead minnow (immature), <i>Pimephales promelas</i>	F, M	Nickel chloride	210	28,000	8,316	-	Pickerling 1974
Fathead minnow (immature), <i>Pimephales promelas</i>	F, M	Nickel chloride	210	25,000	7,425	-	Pickerling 1974
Fathead minnow, <i>Pimephales promelas</i>	F, M	Nickel sulfate	45	5,209	5,695	-	Lind et al. Manuscript
Fathead minnow, <i>Pimephales promelas</i>	F, M	Nickel sulfate	44	5,163	5,753	8,027	Lind et al. Manuscript
Banded killifish (<20 cm), <i>Fundulus diaphanus</i>	S, M	Nickel nitrate	53	46,200†	43,980	-	Rehwoldt et al. 1971
Banded killifish, <i>Fundulus diaphanus</i>	S, M	-	55	46,100	42,530	43,250	Rehwoldt et al. 1972
Guppy (6 mo), <i>Poecilia reticulata</i>	S, U	Nickel chloride	20	4,450	9,661	9,661	Pickerling and Henderson 1966
White perch (<20 cm), <i>Morone americana</i>	S, M	Nickel nitrate	53	13,600†	12,950	-	Rehwoldt et al. 1971
White perch, <i>Morone americana</i>	S, M	-	55	13,700	12,640	12,790	Rehwoldt et al. 1972
Striped bass (fingerling), <i>Morone saxatilis</i>	S, M	Nickel nitrate	53	6,200†	5,902	-	Rehwoldt et al. 1971
Striped bass, <i>Morone saxatilis</i>	S, M	-	55	6,300	5,812	-	Rehwoldt et al. 1972
Striped bass (63 day), <i>Morone saxatilis</i>	S, U	Nickel chloride	40	3,900	4,710	-	Palawski et al. 1985

Table 1. (continued)

<u>Species</u>	<u>Method^a</u>	<u>Chemical</u>	<u>Hardness (mg/L as CaCO₃)</u>	<u>LC50 or EC50 (μg/L)^b</u>	<u>Adjusted LC50 or EC50 (μg/L)^b</u>	<u>Species Mean Acute Value (μg/L)^b</u>	<u>Reference</u>
<u>Striped bass (63 day), <i>Morone saxatilis</i></u>	S, U	Nickel chloride	285	33,000	7,569	5,914	Palawski et al. 1985
<u>Rock bass, <i>Ambloplites rupestris</i></u>	F, M	Nickel sulfate	26	2,480	4,312	4,312	Lind et al. Manuscript
<u>Pumpkinseed (<20 cm), <i>Lepomis gibbosus</i></u>	S, M	Nickel nitrate	53	8,100 ^f	7,710	-	Rahwoldt et al. 1971
<u>Pumpkinseed, <i>Lepomis gibbosus</i></u>	S, M	-	55	8,000	7,380	7,544	Rahwoldt et al. 1972
<u>Bluegill (1-2 g), <i>Lepomis macrochirus</i></u>	S, U	Nickel chloride	20	5,180	11,250	-	Pickering and Henderson 1966
<u>Bluegill (1-2 g), <i>Lepomis macrochirus</i></u>	S, U	Nickel chloride	20	5,360	11,640	-	Pickering and Henderson 1966
<u>Bluegill (1-2 g), <i>Lepomis macrochirus</i></u>	S, U	Nickel chloride	360	39,600	7,454	-	Pickering and Henderson 1966
<u>Bluegill, <i>Lepomis macrochirus</i></u>	F, M	Nickel chloride	49	21,200	21,570	12,040	Cairns et al. 1981

Table 1. (continued)

<u>Species</u>	<u>Method*</u>	<u>Chemical</u>	<u>Salinity (g/kg)</u>	<u>LC50 or EC50 (µg/L)**</u>	<u>Species Mean Acute Value (µg/L)</u>	<u>Reference</u>
<u>Polychaete worm (adult), <i>Nereis arenaceodentata</i></u>	S, U	Nickel chloride	-	49,000	49,000	Petrlich and Reish 1979
<u>Polychaete worm (adult), <i>Nereis virens</i></u>	S, U	Nickel chloride	20	25,000	25,000	Eisler and Hennekey 1977
<u>Polychaete worm (adult), <i>Ctenodrilus serratus</i></u>	S, U	Nickel chloride	-	17,000	17,000	Petrlich and Reish 1979
<u>Polychaete worm (adult), <i>Capitella capitata</i></u>	S, U	Nickel chloride	-	>50,000	>50,000	Petrlich and Reish 1979
<u>Mud snail (adult), <i>Nassarius obsoletus</i></u>	S, U	Nickel chloride	20	72,000	72,000	Eisler and Hennekey 1977
<u>Eastern oyster (embryo), <i>Crassostrea virginica</i></u>	S, U	Nickel chloride	25	1,180	1,180	Calabrese et al. 1975
<u>Clam, <i>Macoma balthica</i></u>	S, U	Nickel chloride	15 (5 °C)	100,000	-	Bryant et al. 1985
<u>Clam, <i>Macoma balthica</i></u>	S, U	Nickel chloride	25 (5 °C)	380,000	-	Bryant et al. 1985
<u>Clam, <i>Macoma balthica</i></u>	S, U	Nickel chloride	35 (5 °C)	700,000	-	Bryant et al. 1985
<u>Clam, <i>Macoma balthica</i></u>	S, U	Nickel chloride	15 (10 °C)	95,000	-	Bryant et al. 1985
<u>Clam, <i>Macoma balthica</i></u>	S, U	Nickel chloride	25 (10 °C)	560,000	-	Bryant et al. 1985
<u>Clam, <i>Macoma balthica</i></u>	S, U	Nickel chloride	35 (10 °C)	1,100,000	-	Bryant et al. 1985
<u>Clam, <i>Macoma balthica</i></u>	S, U	Nickel chloride	15 (15 °C)	110,000	-	Bryant et al. 1985

Table I. (continued)

Species	Method*	Chemical	Salinity (g/kg)	LC50 or EC50 (µg/L)**	Species Mean Acute Value (µg/L)	Reference
Clam, <i>Macoma balthica</i>	S, U	Nickel chloride	25 (15°C)	180,000	-	Bryant et al. 1985
Clam, <i>Macoma balthica</i>	S, U	Nickel chloride	35 (15°C)	540,000	294,500	Bryant et al. 1985
Quahog clam (embryo), <i>Meretrix mercenaria</i>	S, U	Nickel chloride	25	310	-	Calabrese and Nelson 1974
Soft-shell clam (adult), <i>Mya arenaria</i>	S, U	Nickel chloride	20	320,000	-	Eisler and Hennekey 1977
Soft-shell clam (adult), <i>Mya arenaria</i>	S, U	Nickel chloride	30	>50,000	320,000	Eisler 1977a
Copepod (adult), <i>Eurytemora affinis</i>	S, U	Nickel chloride	30	13,180	-	Lussier and Cardin 1985
Copepod (adult), <i>Eurytemora affinis</i>	S, U	Nickel chloride	30	9,593	11,240	Lussier and Cardin 1985
Copepod (adult), <i>Acartia clausi</i>	S, U	Nickel chloride	30	3,466	3,466	Lussier and Cardin 1985
Copepod (adult), <i>Nitocra spinipes</i>	S, U	Nickel chloride	7	6,000	6,000	Bengtsson 1978
Mysid (juvenile), <i>Icteromyces formosa</i>	S, M	Nickel chloride	30	151.7	151.7	Gentile et al. 1982
Mysid (juvenile), <i>Mysidopsis bahia</i>	F, M	Nickel chloride	30	508	508	Gentile et al. 1982; Lussier et al. 1985
Mysid (juvenile), <i>Mysidopsis bigelowi</i>	S, M	Nickel chloride	30	634	634	Gentile et al. 1982
Amphipod, <i>Corophium volutator</i>	S, U	Nickel chloride	5 (5°C)	5,000	-	Bryant et al. 1985
Amphipod, <i>Corophium volutator</i>	S, U	Nickel chloride	10 (5°C)	21,000	-	Bryant et al. 1985

Table 1. (continued)

Species	Method*	Chemical	Salinity (g/kg)	LC50 or EC50 ($\mu\text{g/L}$)**	Species Mean Acute Value ($\mu\text{g/L}$)	Reference
Amphipod, <i>Corophium volutator</i>	S, U	Nickel chloride	15 (5°C)	18,000	-	Bryant et al. 1985
Amphipod, <i>Corophium volutator</i>	S, U	Nickel chloride	25 (5°C)	36,000	-	Bryant et al. 1985
Amphipod, <i>Corophium volutator</i>	S, U	Nickel chloride	35 (5°C)	54,000	-	Bryant et al. 1985
Amphipod, <i>Corophium volutator</i>	S, U	Nickel chloride	5 (10°C)	9,000	-	Bryant et al. 1985
Amphipod, <i>Corophium volutator</i>	S, U	Nickel chloride	10 (10°C)	15,000	-	Bryant et al. 1985
Amphipod, <i>Corophium volutator</i>	S, U	Nickel chloride	15 (10°C)	22,000	-	Bryant et al. 1985
Amphipod, <i>Corophium volutator</i>	S, U	Nickel chloride	25 (10°C)	24,000	-	Bryant et al. 1985
Amphipod, <i>Corophium volutator</i>	S, U	Nickel chloride	35 (10°C)	52,000	-	Bryant et al. 1985
Amphipod, <i>Corophium volutator</i>	S, U	Nickel chloride	5 (15°C)	5,600	-	Bryant et al. 1985
Amphipod, <i>Corophium volutator</i>	S, U	Nickel chloride	10 (15°C)	16,000	-	Bryant et al. 1985
Amphipod, <i>Corophium volutator</i>	S, U	Nickel chloride	15 (15°C)	18,000	-	Bryant et al. 1985
Amphipod, <i>Corophium volutator</i>	S, U	Nickel chloride	25 (15°C)	22,000	-	Bryant et al. 1985
Amphipod, <i>Corophium volutator</i>	S, U	Nickel chloride	35 (15°C)	34,000	18,950	Bryant et al. 1985
Hermit crab (adult), <i>Pagurus longicarpus</i>	S, U	Nickel chloride	20	47,000	47,000	Eisler and Hennekey 1977

Table 1. (continued)

<u>Species</u>	<u>Method*</u>	<u>Chemical</u>	<u>Salinity (g/kg)</u>	<u>LC50 or EC50 (µg/L)**</u>	<u>Species Mean Acute Value (µg/L)</u>	<u>Reference</u>
Starfish (adult), <i>Asterias forbesii</i>	S, U	Nickel chloride	20	150,000	150,000	Elsler and Hennaker 1977
Mumm Ichog (adult), <i>Fundulus heteroclitus</i>	S, U	Nickel chloride	6.9	55,000	-	Dorfman 1977
Mumm Ichog (adult), <i>Fundulus heteroclitus</i>	S, U	Nickel chloride	21.6	175,000	-	Dorfman 1977
Mumm Ichog (adult), <i>Fundulus heteroclitus</i>	S, U	Nickel chloride	20	350,000	149,900	Elsler and Hennaker 1977
Atlantic silverside (larva), <i>Menidia menidia</i>	S, U	Nickel chloride	30	7,958	7,958	Cardin 1985
Tidewater silverside (juvenile), <i>Menidia peninsulae</i>	S, U	Nickel chloride	20	38,000	38,000	Hansen 1983
Striped bass (65 day), <i>Morone saxatilis</i>	S, U	Nickel chloride	1	21,000	21,000	Palawski et al. 1985
Spot (juvenile), <i>Lepostomus xanthurus</i>	S, U	Nickel chloride	21	70,000	70,000	Hansen 1983

* S = static; R = renewal; F = flow-through; M = measured; U = unmeasured.

** Results are expressed as nickel, not as the chemical.

*** Freshwater LC50s and EC50s were adjusted to hardness = 50 mg/l (as CaCO₃) using the pooled slope of 0.8460 (see text).

**** Freshwater Species Mean Acute Values are calculated at hardness = 50 mg/l (as CaCO₃).

† In river water.

Table 1. (continued)

Results of Covariance Analysis of Freshwater Acute Toxicity versus Hardness

<u>Species</u>	<u>n</u>	<u>Slope</u>	<u>95% Confidence Limits</u>	<u>Degrees of Freedom</u>
<u>Daphnia magna</u>	6	1.1810	0.3187, 2.0433	4
<u>Fathead minnow</u>	10	0.8294	0.6755, 0.9833	8
<u>Striped bass</u>	4	1.0459	0.7874, 1.3045	2
<u>Bluegill</u>	4	0.6909	-0.1654, 1.5472	2
All of above	24	0.8460*	0.7004, 0.9915	19

* P = 0.26 for equality of slopes with 16 degrees of freedom.

Table 2. Chronic Toxicity of Nickel to Aquatic Animals

<u>Species</u>	<u>Test*</u>	<u>Cheical</u>	<u>FRESHWATER SPECIES</u>	<u>Hardness (mg/L as CaCO₃)</u>	<u>Limits (µg/L)^{**}</u>	<u>Chronic Value (µg/L)</u>	<u>Reference</u>
Cladoceran, <u>Daphnia magna</u>	LC	Nickel chloride	51	10.2- 21.4	14.77	Chapman et al. Manuscript	
Cladoceran, <u>Daphnia magna</u>	LC	Nickel chloride	105	101- 150	123.1	Chapman et al. Manuscript	
Cladoceran, <u>Daphnia magna</u>	LC	Nickel chloride	205	220- 578	356.6	Chapman et al. Manuscript	
Cladoceran (1st generation), <u>Daphnia magna</u>	LC	Nickel sulfate	-	5-10	7.071	Lazareva 1985	
Cladoceran (2nd generation), <u>Daphnia magna</u>	LC	Nickel sulfate	-	<5***	<5	Lazareva 1985	
Cladoceran (3rd generation), <u>Daphnia magna</u>	LC	Nickel sulfate	-	5-10	7.071	Lazareva 1985	
Cladoceran (4th generation), <u>Daphnia magna</u>	LC	Nickel sulfate	-	5-10	7.071	Lazareva 1985	
Caddisfly, <u>Clistoronia magnifica</u>	LC	Nickel chloride	54	66- 250	128.4	Nebeker et al. 1984	
Rainbow trout, <u>Salmo gairdneri</u>	ELS	Nickel chloride	53	<35***	<35	Nebeker et al. 1985	
Rainbow trout, <u>Salmo gairdneri</u>	ELS	Nickel chloride	52	62- 134	91.15	Nebeker et al. 1985	
Rainbow trout, <u>Salmo gairdneri</u>	ELS	Nickel chloride	49	134- 431	240.3	Nebeker et al. 1985	
Fatherhead minnow, <u>Pimephales promelas</u>	LC	Nickel chloride	210	380- 730	526.7	Pickering 1974	
Fatherhead minnow, <u>Pimephales promelas</u>	ELS	Nickel sulfate	44† 45†	108.9- 433.5	217.3	Lind et al. Manuscript	

Table 2. (Continued)

<u>Species</u>	<u>Test^a</u>	<u>Chemical</u>	<u>Salinity (g/kg)</u>	<u>Limits (µg/L)^b</u>	<u>Chronic Value (µg/L)</u>	<u>Reference</u>
		SALTWATER SPECIES				
Mysid, <i>Mysidopsis bahia</i>	LC	Nickel chloride	30	61- 141	92.74	Gentile et al. 1982; Lussier et al. 1985

* LC = life-cycle or partial life-cycle; ELS = early life-stage.

** Results are based on measured concentrations of nickel.

*** Unacceptable effects occurred at all concentrations tested.

† Values from acute tests in Table 1.

Results of Regression Analysis of Freshwater Chronic Toxicity versus Hardness

<u>Species</u>	<u>n</u>	<u>Slope</u>	<u>95% Confidence Limits</u>	<u>Degrees of Freedom</u>
<u>Daphnia magna</u>	3	2.3007	-2.6551, 7.2568	1
Fathead minnow	2	0.5706	*	0
All of above	5	1.3418**	-1.3922, 4.0760	2

* Cannot be calculated because degrees of freedom = 0.

** P = 0.19 for equality of slopes with 1 degree of freedom.

Table 2. (Continued)

Acute-Chronic Ratio

<u>Species</u>	<u>Hardness (mg/L as CaCO₃)</u>	<u>Acute Value (µg/L)</u>	<u>Chronic Value (µg/L)</u>	<u>Ratio</u>
Cladoceran, <u>Daphnia magna</u>	51	1,800	14.77	122.4
Cladoceran, <u>Daphnia magna</u>	104- 105	1,920	123.1	15.60
Cladoceran, <u>Daphnia magna</u>	205- 206	4,970	356.6	13.94
Fathead minnow, <u>Pimephales promelas</u>	210	27,930*	526.7	53.03
Fathead minnow, <u>Pimephales promelas</u>	44- 45	5,186**	217.3	23.87
Mysid, <u>Mysidopsis bahia</u>	30***	508	92.74	5.478

* Geometric mean of four values in Table 1.

** Geometric mean of two values in Table 1.

*** Salinity (g/kg).

Table 3. Ranked Genus Mean Acute Values with Species Mean Acute-Chronic Ratios

Rank #	Genus Mean Acute Value ($\mu\text{g/L}$)	Species	FRESHWATER SPECIES		
			Species Mean Acute Value ($\mu\text{g/L}$)	Mean Acute-Chronic Ratio	Mean Acute-Chronic Ratio
18	43,250	Banded killifish, <u>Fundulus diaphanus</u>	43,250	-	-
17	40,460	Stonefly, <u>Acroneuria lycorias</u>	40,460	-	-
16	30,200	Caddisfly, Unidentified sp.	30,200	-	-
15	21,320	Goldfish, <u>Carassius auratus</u>	21,320	-	-
14	21,200	Damsel fly, Unidentified sp.	21,200	-	-
13	14,100	Worm, <u>Nais</u> sp.	14,100	-	-
12	13,380	Rainbow trout, <u>Salmo gairdneri</u>	13,380	-	-
11	13,000	Amphipod, <u>Gammarus</u> sp.	13,000	-	-
10	12,770	Snail, <u>Amnicola</u> sp.	12,770	-	-
9	12,180	American eel, <u>Anguilla rostrata</u>	12,180	-	-
8	9,839	Common carp, <u>Cyprinus carpio</u>	9,839	-	-
7	9,661	Guppy, <u>Poecilia reticulata</u>	9,661	-	-
6	8,697	White perch, <u>Morone americana</u>	12,790	-	-
		Striped bass, <u>Morone saxatilis</u>	5,914	-	-

Table 3. (Continued)

<u>Rank*</u>	<u>Genus Mean Acute Value (μg/L)†</u>	<u>Species</u>	<u>Species Mean Acute Value (μg/L)‡</u>	<u>Species Mean Acute-Chronic Ratio***</u>
5	9,530	Pumpkinseed, <u>Lepomis gibbosus</u>	7,544	-
4	6,027	Bluegill, <u>Lepomis macrochirus</u>	12,040	-
3	4,636	Fathead minnow, <u>Pimephales promelas</u>	8,027	35.58†
2	4,312	Mayfly, <u>Ephemerella subvaria</u>	4,636	-
1	1,500	Rock bass, <u>Ambloplites rupestris</u>	4,312	-
		Cladoceran, <u>Daphnia pulex</u>	2,042	-
		Cladoceran, <u>Daphnia magna</u>	1,102	29.86††
<u>SALTWATER SPECIES</u>				
20	320,000	Soft-shell clam, <u>Mya arenaria</u>	320,000	-
19	294,500	Clam, <u>Macoma balthica</u>	294,500	-
18	150,000	Starfish, <u>Asterias forbesii</u>	150,000	-
17	149,900	Mummichog, <u>Fundulus heteroclitus</u>	149,900	-
16	72,000	Mud snail, <u>Nassarius obsoletus</u>	72,000	-
15	70,000	Spot, <u>Leiostomus xanthurus</u>	70,000	-

Table 3. (Cont'd)

Rank#	Genus Mean Acute Value ($\mu\text{g/L}$) ^a	Species	Species Mean Acute Value ($\mu\text{g/L}$) ^a	Species Mean Acute-Chronic Ratio ^b
14	>50,000	Polychaete worm, <u>Capitella capitata</u>	>50,000	-
13	47,000	Hermit crab, <u>Pagurus longicarpus</u>	47,000	-
12	35,000	Polychaete worm, <u>Nereis arenaceodentata</u>	49,000	-
		Polychaete worm, <u>Nereis virens</u>	25,000	-
11	21,000	Striped bass, <u>Morone saxatilis</u>	21,000	-
10	17,390	Atlantic silverside, <u>Menidia menidia</u>	7,958	-
		Tidewater silverside, <u>Menidia peninsulae</u>	38,000	-
9	17,000	Polychaete worm, <u>Ctenodrilus serratus</u>	17,000	-
8	18,950	Amphipod, <u>Corophium volutator</u>	18,950	-
7	11,240	Copepod, <u>Eurytemora affinis</u>	11,240	-
6	6,000	Copepod, <u>Nitocra spinipes</u>	6,000	-
5	3,466	Copepod, <u>Acartia clausi</u>	3,466	-
4	1,180	Eastern oyster, <u>Crassostrea virginica</u>	1,180	-

Table 3. (Continued)

<u>Rank*</u>	<u>Genus Mean Acute Value (µg/L)†</u>	<u>Species</u>	<u>Species Mean Acute Value (µg/L)***</u>	<u>Species Mean Acute-Chronic Ratios**</u>
3	567.5	Mysid, <u>Mysidopsis bahia</u>	508	5.476
2	310	Mysid, <u>Mysidopsis bigelowi</u>	634	-
1	151.7	Quahog clam, <u>Meretrix mercenaria</u>	310	-
		Heteronyx is formosa	151.7	-

* Ranked from most resistant to most sensitive based on Genus Mean Acute Value. Inclusion of "greater than" values does not necessarily imply a true ranking, but does allow use of all genera for which data are available so that the Final Acute Value is not unnecessarily lowered.

** Freshwater Genus Mean Acute Values are at hardness = 50 mg/L.

*** From Table 1; freshwater values are at hardness = 50 mg/L.

**** From Table 2.

† Geometric mean of two values in Table 2.

** Geometric mean of three values in Table 2.

Table 3. (Continued)

<u>Fresh water</u>
Final Acute Value = 1,578 µg/L (at hardness = 50 mg/L)
Cr1terion Maximum Concentration = (1,578 µg/L) / 2 = 789.0 µg/L (at hardness = 50 mg/L)
Pooled Slope = 0.8460 (see Table 1)
$\ln(\text{Criterion Maximum Concentration}) = \ln(789.0) - [\text{slope} \times \ln(50)]$
$= 6.6708 - (0.8460 \times 3.9120) = 3.3612$
Criterion Maximum Concentration = $e^{(0.8460[\ln(\text{hardness})] + 3.3612)}$
Final Acute-Chronic Ratio = 17.99 (see text)
Final Chronic Value = (1,578 µg/L) / 17.99 = 87.72 µg/L (at hardness = 50 mg/L)
Assumed Chronic Slope = 0.8460 (see text)
$\ln(\text{Final Chronic Concentration}) = \ln(87.72) - [\text{slope} \times \ln(50)]$
$= 4.4741 - (0.8460 \times 3.9120) = 1.1645$
Final Chronic Value = $e^{(0.8460[\ln(\text{hardness})] + 1.1645)}$
<u>Salt water</u>
Final Acute Value = 149.2 µg/L
Criterion Maximum Concentration = 149.2 / 2 = 74.60 µg/L
Final Acute-Chronic Ratio = 17.99 (see text)
Final Chronic Value = (149.2 µg/L) / 17.99 = 8.295 µg/L

Table 4. Toxicity of Nickel to Aquatic Plants

<u>Species</u>	<u>Chemical</u>	<u>Hardness (mg/L as CaCO_3)</u>	<u>Duration (days)</u>	<u>Effect</u>	<u>Concentration ($\mu\text{g/L}$)^a</u>	<u>Reference</u>
<u>FRESHWATER SPECIES</u>						
Blue-green alga, <i>Anabaena flos-aquae</i>	Nickel nitrate	-	14	84% reduction In growth	600	Spencer and Greene 1981
Blue-green alga, <i>Microcystis aeruginosa</i>	Nickel chloride	-	8	Incipient inhibition	5	Bringmann and Kuhn 1978a,b
Green alga, <i>Ankistrodesmus falcatus</i>	Nickel chloride	-	10	45% reduction In growth	5,000	Dev I Prasad and Dev I Prasad 1982
Green alga, <i>Ankistrodesmus falcatus</i>	Nickel nitrate	-	14	98% reduction In growth	100	Spencer and Greene 1981
Green alga, <i>Ankistrodesmus falcatus</i> var. <u>acicularis</u>	Nickel nitrate	-	14	42% reduction In growth	100	Spencer and Greene 1981
Green alga, <i>Chlamydomonas eugametos</i>	Nickel nitrate or Nickel sulfate	47.5	12	91% reduction In growth	700**	Hutchinson 1973; Hutchinson and Stokes 1975
Green alga, <i>Chlorella vulgaris</i>	Nickel nitrate or Nickel sulfate	47.5	12	53% reduction In growth	300**	Hutchinson 1973; Hutchinson and Stokes 1975
Green alga, <i>Chlorococcum</i> sp.	Nickel chloride	-	10	52% reduction In growth	5,000	Dev I Prasad and Dev I Prasad 1982
Green alga, <i>Haematococcus capensis</i>	Nickel nitrate or Nickel sulfate	47.5	12	85% reduction In growth	300**	Hutchinson 1973; Hutchinson and Stokes 1975
Green alga, <i>Pedialastrum tetrads</i>	Nickel nitrate	-	14	Increased growth	100	Spencer and Greene 1981
Green alga, <i>Scenedesmus acuminata</i>	Nickel nitrate or Nickel sulfate	47.5	12	54% reduction In growth	50**	Hutchinson 1973; Hutchinson and Stokes 1975

Table 4. (Continued)

<u>Species</u>	<u>Chemical</u>	<u>Hardness (mg/L as CaCO_3)</u>	<u>DURATION (days)</u>	<u>Effect</u>	<u>Concentration ($\mu\text{g/L}$)^a</u>	<u>Reference</u>
<u>Green alga, <i>Scenedesmus acuminata</i></u>	Nickel nitrate or Nickel sulfate	47.5	13	Reduced growth	500	Stokes et al. 1973; Hutchinson and Stokes 1975
<u>Green alga, <i>Scenedesmus dimorphus</i></u>	Nickel nitrate	-	14	30% reduction in growth	100	Spencer and Greene 1981
<u>Green alga, <i>Scenedesmus obliquus</i></u>	Nickel chloride	-	10	47% reduction in growth	3,000	Dev Prasad and Dev Prasad 1982
<u>Green alga, <i>Scenedesmus quadrivalvis</i></u>	Nickel chloride	-	8	Incomplete inhibition	1,300	Berlingmann and Kuhn 1977a; 1978a,b; 1979; 1980b
<u>Green alga, <i>Scenedesmus quadrivalvis</i></u>	Nickel nitrate	-	14	60% reduction in growth	100	Spencer and Greene 1981
<u>Diatom, <i>Navicula pelluculosa</i></u>	Nickel nitrate	14.96	14	82% reduction in growth	100	Fazy et al. 1979
<u>Duckweed, <i>Lemna minor</i></u>	Nickel chloride	-	28	EC50	340	Brown and Rattigan 1979
<u>Duckweed, <i>Lemna minor</i></u>	-	-	4	EC50 (growth)	450	Wang 1986
<u>Macrophyte, <i>Eloea (Anacharis) canadensis</i></u>	Nickel chloride	-	28	EC50	2,800	Brown and Rattigan 1979
<u>Water hyacinth, <i>Elodea crassipes</i></u>	Nickel chloride	12	38	30% reduction in growth	4,000	Muramoto and Okita 1984
<u>Water hyacinth, <i>Elodea crassipes</i></u>	Nickel chloride	12	38	29% reduction in growth	8,000	Muramoto and Okita 1984
<u>SALTWATER SPECIES</u>						
<u>Giant kelp (young fronds), <i>Macrocystis pyrifera</i></u>	-	4	-	EC50 (reduc- tion in photosynthesis)	2,000	Cleland and North 1959

* Results are expressed as nickel, not as the chemical.

Table 5. Bioaccumulation of Nickel by Aquatic Organisms

<u>Species</u>	<u>Chemical</u>	<u>Concentration in Water ($\mu\text{g/L}$)^a</u>	<u>Hardness (mg/L as CaCO_3)</u>	<u>Duration (days)</u>	<u>Tissue</u>	<u>BCF or BAF^b</u>	<u>Reference</u>
FRESHWATER SPECIES							
Green alga, <i>Scenedesmus acuminata</i>	Nickel nitrate or Nickel sulfate	1,000	-	6	Whole body	9.3	Hutchinson and Stokes 1975
Water hyacinth, <i>Eichornia crassipes</i>	Nickel chloride	1,000	38	12	Root tops	256.0 174.2	Muramoto and Oki 1984
Water hyacinth, <i>Eichornia crassipes</i>	Nickel chloride	4,000	38	12	Root tops	438.2 500.3	Muramoto and Oki 1984
Water hyacinth, <i>Eichornia crassipes</i>	Nickel chloride	8,000	38	12	Root tops	335.5 576.2	Muramoto and Oki 1984
Cladoceran, <i>Daphnia magna</i>	^{63}Ni in 0.1M HCl	-	-	-	Whole body	100	Hall 1978
Cladoceran, <i>Daphnia magna</i>	-	50	20.1	3.75	Whole body	192****	Hall 1982
Cladoceran, <i>Daphnia magna</i>	-	750	20.1	3.75	Whole body	123***	Hall 1982
Cladoceran, <i>Daphnia magna</i>	Nickel chloride	58.7	-	13	Whole body	11.6	Watras et al. 1985
Cladoceran, <i>Daphnia magna</i>	Nickel chloride	1,000	320	180	Muscle	0.8	Calamar et al. 1982
Rainbow trout, <i>Salmo gairdneri</i>	Nickel sulfate	-	-	30	Whole body	106	Lind et al. Manuscript
Fathead minnow, <i>Pimephales promelas</i>	Nickel sulfate	21	-	-	Whole body	79	Lind et al. Manuscript
Fathead minnow, <i>Pimephales promelas</i>	Nickel sulfate	44.4	-	30	Whole body	-	Lind et al. Manuscript
Fathead minnow, <i>Pimephales promelas</i>	Nickel sulfate	108.9	-	30	Whole body	47	Lind et al. Manuscript
SALTWATER SPECIES							
Rockweed, <i>Fucus vesiculosus</i>	-	1.2	-	Field collections	Whole plant	675†	Foster 1976

Table 5. (Continued)

<u>Species</u>	<u>Chemical</u>	<u>Concentration In Water (µg/L)*</u>	<u>Hardness (mg/L as CaCO_3)*</u>	<u>Duration (days)</u>	<u>Tissue</u>	<u>BCF or BAFF**</u>	<u>Reference</u>
Brown macroalgae, <i>Ascophyllum nodosum</i>	-	1.2	-	Field collections	whole plant	458.3†	Foster 1976
Blue mussel, <i>Mytilus edulis</i>	Nickel sulfate	4.4	-	84	Soft parts	472.7	Zarogian and Johnson 1984
Blue mussel, <i>Mytilus edulis</i>	Nickel sulfate	10.0	-	84	Soft parts	328.6	Zarogian and Johnson 1984
Eastern oyster, <i>Crassostrea virginica</i>	Nickel sulfate	4.2	-	84	Soft parts	458.1	Zarogian and Johnson 1984
Eastern oyster, <i>Crassostrea virginica</i>	Nickel sulfate	9.9	-	84	Soft parts	261.8	Zarogian and Johnson 1984

* Measured concentration of nickel.

** Bioconcentration factors (BCFs) and bioaccumulation factors (BAFs) are based on measured concentrations of nickel in water and in tissue.

*** Estimated from graph.

† Factor was converted from dry weight to wet weight basis.

Table 6. Other Data on Effects of Nickel on Aquatic Organisms

<u>Species</u>	<u>Chemical</u>	<u>Hardness (mg/L as CaCO_3)</u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration ($\mu\text{g/L}$)^a</u>	<u>Reference</u>
<u>FRESHWATER SPECIES</u>						
Alga, <i>Chlorella pyrenoidosa</i>	-	-	24 hr	Reduced growth	88	Gerhards and Weller 1977
Green alga, <i>Scenedesmus quadrivalva</i>	Nickel chloride	-	96 hr	Inhibition (river water)	1,500	Brlingmann and Kuhn 1959a,b
Green alga, <i>Scenedesmus quadrivalva</i>	Nickel ammonium sulfate	-	96 hr	Inhibition (river water)	900	Brlingmann and Kuhn 1959a,b
Alga, (mixed population)	Nickel nitrate	87-99	<53 days	Decrease in diversity; shift to green and blue-green algae	2-8.6	Patrick et al. 1975
Blue-green alga, <i>Anabaena cylindrica</i>	Nickel sulfate	-	5 days	No effect on doubling time (in light)	15.1	Daday et al. 1985
Blue-green alga, <i>Anabaena cylindrica</i>	Nickel sulfate	-	5 days	13% reduction in doubling time (in dark)	15.1	Daday et al. 1985
Blue-green alga, <i>Anabaena cylindrica</i>	Nickel chloride	-	30 hr	BCF = 680.5 (in light)	-	Campbell and Smith 1986
Blue-green alga, <i>Anabaena cylindrica</i>	Nickel chloride	-	30 hr	BCF = 375.0 (in dark)	-	Campbell and Smith 1986
Blue-green alga, <i>Nostoc linckia</i>	Nickel chloride	-	24 hr	EC50 (nitrate reduction)	1,885	Kumar et al. 1985
Blue-green alga, <i>Nostoc linckia</i>	Nickel chloride	-	24 hr	EC50 (ammonia uptake)	1,141	Kumar et al. 1985
Blue-green alga, <i>Nostoc muscorum</i>	Nickel chloride	-	21 days	EC50 (survival)	235.1	Rai and Ralzada 1985
Bacterium, <i>Aeromonas sobria</i>	Nickel chloride	40	24 hr**	Reduction in abundance	5	Bablich and Stotzky 1983

Table 6. (continued)

<u>Species</u>	<u>Chemical</u>	<u>Hardness (mg/L as CaCO₃)</u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration (µg/L)^a</u>	<u>Reference</u>
<i>Bacterium, Bacillus brevis</i>	Nickel chloride	40	24 hr**	Reduction in abundance	5	Bablich and Stotzky 1983
<i>Bacterium, Bacillus cereus</i>	Nickel chloride	40	24 hr**	Reduction in abundance	5	Bablich and Stotzky 1983
<i>Bacterium, Escherichia coli</i>	Nickel chloride	-	-	Inhibition	100	Brüngmann and Kuhn 1959a
<i>Bacterium, Escherichia coli</i>	Nickel ammonium sulfate	-	-	Inhibition	100	Brüngmann and Kuhn 1959a
<i>Bacterium, Pseudomonas putida</i>	Nickel chloride	-	16 hr	Inhibition	2.5 (3.0)	Brüngmann and Kuhn 1977a; 1979; 1980b
<i>Bacterium, Serratia marcescens</i>	Nickel chloride	40	24 hr**	Reduction in abundance	10	Bablich and Stotzky 1983
<i>Bacterium, Nitrosomonas europaea</i>	-	9	-	No growth	400	Sato et al. 1986
<i>Mixed heterotrophic bacteria</i>	Nickel chloride	-	0.5 hr	EC50 (survival)	42.9	Arlbright et al. 1972
<i>Protozoan, Entosiphon sulcatum</i>	Nickel chloride	-	72 hr	Inhibition	140	Brüngmann 1978; Brüngmann and Kuhn 1979; 1980b; 1981
<i>Protozoan, Microregma heterostoma</i>	Nickel chloride	-	28 hr	Inhibition	50	Brüngmann and Kuhn 1959b
<i>Protozoan, Microregma heterostoma</i>	Nickel ammonium sulfate	-	28 hr	Inhibition	70	Brüngmann and Kuhn 1959b
<i>Protozoan, Chilomonas paramecium</i>	Nickel chloride	-	48 hr	Inhibition	820	Brüngmann et al. 1980; Brüngmann and Kuhn 1981
<i>Protozoan, Uronema parduezi</i>	Nickel chloride	-	20 hr	Inhibition	42	Brüngmann and Kuhn 1980a, 1981

Table 6. (Continued)

<u>Species</u>	<u>Chemical</u>	<u>Hardness (mg/l as CaCO_3)</u>	<u>Durat ion</u>	<u>Effect</u>	<u>Concentration ($\mu\text{g/L}$)^a</u>	<u>Reference</u>
Tubifex worm, Tubifex tubifex	Nickel sulfate	34.2	48 hr	LC50	8.70 7.00	Brekov Ic-Popov Ic and Popov Ic 1977a
Cladoceran, <u>Daphnia magna</u>	Nickel chloride	-	48 hr	EC50 (river water)	6,000	Br Ingmann and Kuhn 1959a,b
Cladoceran, <u>Daphnia magna</u>	Nickel ammonium sulfate	-	48 hr	EC50 (river water)	6,000	Br Ingmann and Kuhn 1959a,b
Cladoceran, <u>Daphnia magna</u>	Nickel chloride	288	24 hr	EC50 (swimming)	11,000	Br Ingmann and Kuhn 1977b
Cladoceran, <u>Daphnia magna</u>	Nickel chloride	45.3	48 hr	EC50 (immobi- lization) (fed)	1,120	Blesinger and Christensen 1972
Cladoceran, <u>Daphnia magna</u>	Nickel chloride	45.3	21 days	EC50 (immobi- lization)	130	Blesinger and Christensen 1972
Cladoceran, <u>Daphnia magna</u>	Nickel chloride	45.3	21 days	16% reproduc- tive impairment	30	Blesinger and Christensen 1972
Cladoceran, <u>Daphnia magna</u>	Nickel chloride	-	72 hr	BCF = 0.823 BCF = 0.526 BCF = 1.83 BCF = 2.20 BCF = 1.17	1,855 1,115 185.5 58.70 18.50	Watras et al. 1985
Cladoceran, <u>Daphnia magna</u>	Nickel sulfate	25	48 hr	LC50 (TOC = 39 mg/L)	2,171	Lind et al. Manuscript
Cladoceran, <u>Daphnia pulex</u>	Nickel sulfate	28	48 hr	LC50 (TOC = 15 mg/L)	1,140	Lind et al. Manuscript
Cladoceran, <u>Daphnia pulex</u>	Nickel sulfate	28	48 hr	LC50 (TOC = 13 mg/L)	1,034	Lind et al. Manuscript
Cladoceran, <u>Daphnia pulex</u>	Nickel sulfate	29	48 hr	LC50 (TOC = 13 mg/L)	697	Lind et al. Manuscript

Table 6. (Continued)

<u>Species</u>	<u>Chemical</u>	<u>Hardness (mg/L as CaCO_3)</u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration ($\mu\text{g/L}$)^a</u>	<u>Reference</u>
Cladoceran, <u>Daphnia pulicaria</u>	Nickel sulfate	73	48 hr	LC50 (TOC = 28 mg/L)	3,414	Lind et al. Manuscript
Cladoceran, <u>Daphnia pulicaria</u>	Nickel sulfate	74	48 hr	LC50 (TOC = 28 mg/L)	2,325	Lind et al. Manuscript
Cladoceran, <u>Daphnia pulicaria</u>	Nickel sulfate	84	48 hr	LC50 (TOC = 32 mg/L)	3,014	Lind et al. Manuscript
Cladoceran, <u>Daphnia pulicaria</u>	Nickel sulfate	86	48 hr	LC50 (TOC = 34 mg/L)	3,316	Lind et al. Manuscript
Cladoceran, <u>Daphnia pulicaria</u>	Nickel sulfate	89	48 hr	LC50 (TOC = 18 mg/L)	2,042	Lind et al. Manuscript
Cladoceran, <u>Daphnia pulicaria</u>	Nickel sulfate	89	48 hr	LC50 (TOC = 34 mg/L)	2,717	Lind et al. Manuscript
Cladoceran, <u>Daphnia pulicaria</u>	Nickel sulfate	100	48 hr	LC50 (TOC = 34 mg/L)	3,757	Lind et al. Manuscript
Cladoceran, <u>Daphnia pulicaria</u>	Nickel sulfate	114	48 hr	LC50 (TOC = 27 mg/L)	3,156	Lind et al. Manuscript
Cladoceran, <u>Daphnia pulicaria</u>	Nickel sulfate	120	48 hr	LC50 (TOC = 33 mg/L)	3,607	Lind et al. Manuscript
Midge, <u>Chironomus</u> sp.	-	50	96 hr	LC50	8,600	Rehwoldt et al. 1975
Coho salmon (yearling), <u>Oncorhynchus kisutch</u>	Nickel chloride	90	144 hr	100% survival	5,000	Lorz et al. 1978
Rainbow trout (0.5-0.9 g), <u>Salmo gairdneri</u>	Nickel sulfate	42	48 hr	LC50	35,750	Willford 1966
Rainbow trout (1 yr), <u>Salmo gairdneri</u>	Nickel sulfate	240	48 hr	LC50	32,000	Brown and Dalton 1970

Table 6. (Continued)

<u>Species</u>	<u>Chemical</u>	<u>Hardness (mg/L as CaCO₃)</u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration (µg/L)^a</u>	<u>Reference</u>
Rainbow trout (embryo, larva), <i>Salmo gairdneri</i>	Nickel chloride	104 (92-110)	28 days	EC50 (death and deformity)	50	Birge 1978; Birge and Black 1980; Birge et al. 1978, 1979, 1980, 1981
Rainbow trout (embryo, larva), <i>Salmo gairdneri</i>	Nickel chloride	125	28 days	EC50 (death and deformity)	60	Birge et al. 1981
Rainbow trout (embryo, larva), <i>Salmo gairdneri</i>	Nickel chloride	174	28 days	EC50 (death and deformity)	90	Birge et al. 1981
Rainbow trout, <i>Salmo gairdneri</i>	Nickel sulfate	240	3.5 days	Decreased gill diffusion	2,000	Hughes et al. 1979
Rainbow trout (adult), <i>Salmo gairdneri</i>	Nickel chloride	320	6 mo	Increase in liver proteolytic activity of males	1,000	Arillo et al. 1982
Rainbow trout (10 g), <i>Salmo gairdneri</i>	Nickel chloride	28.4	20 min threshold	Avoidance threshold	23.9	Giattina et al. 1982
Rainbow trout, <i>Salmo gairdneri</i>	Nickel sulfate	22.5	48 hr	LC50	54,963	Bornatowicz 1983
Rainbow trout (5 days post hatch), <i>Salmo gairdneri</i>	Nickel chloride	50	38 days	LC50	1,400	Nebeker et al. 1985
Brown trout (0.8-1.2 g), <i>Salmo trutta</i>	Nickel sulfate	42	48 hr	LC50	60,290	Willford 1966
Brook trout (0.4-0.6 g), <i>Salvelinus fontinalis</i>	Nickel sulfate	42	48 hr	LC50	54,040	Willford 1966
Lake trout (2.5-3.2 g), <i>Salvelinus namaycush</i>	Nickel sulfate	42	48 hr	LC50	16,750	Willford 1966

Table 6. (Continued)

<u>Species</u>	<u>Chemical</u>	<u>Hardness (mg/L as CaCO_3)</u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration ($\mu\text{g/L}$)^a</u>	<u>Reference</u>
<u>Goldfish, <i>Carassius auratus</i></u>	Nickel chloride	-	19-50 hr 200-210 hr	LT LT	100,000 10,000	Ellis 1957
<u>Goldfish (embryo, larva), <i>Carassius auratus</i></u>	Nickel chloride	195	7 days	EC50 (death and deformity)	2,140	Birge 1978
<u>Goldfish (embryo, larva), <i>Carassius auratus</i></u>	Nickel chloride	93- 105	7 days	EC50 (death and deformity)	2,780	Birge and Black 1980; Birge et al. 1981
<u>Common carp (embryo), <i>Cyprinus carpio</i></u>	Nickel sulfate	128	72 hr	LC50	6,100	Blaylock and Frank 1979
<u>Common carp (larva), <i>Cyprinus carpio</i></u>	Nickel sulfate	128	72 hr 257 hr	LC50	8,460 750	Blaylock and Frank 1979
<u>Common carp (embryo), <i>Cyprinus carpio</i></u>	Nickel sulfate	360	-	EC50 (hatch)	22,000	Kapur and Yadav 1982
<u>Fathead minnow, <i>Pimephales promelas</i></u>	Nickel sulfate	28	96 hr	LC50 (TOC = 14 mg/L)	2,923	Lind et al. Manuscript
<u>Fathead minnow, <i>Pimephales promelas</i></u>	Nickel sulfate	29	96 hr	LC50 (TOC = 12 mg/L)	2,916	Lind et al. Manuscript
<u>Fathead minnow, <i>Pimephales promelas</i></u>	Nickel sulfate	77	96 hr	LC50 (TOC = 32 mg/L)	12,356	Lind et al. Manuscript
<u>Fathead minnow, <i>Pimephales promelas</i></u>	Nickel sulfate	86	96 hr	LC50 (TOC = 15 mg/L)	5,385	Lind et al. Manuscript
<u>Fathead minnow, <i>Pimephales promelas</i></u>	Nickel sulfate	89	96 hr	LC50 (TOC = 33 mg/L)	17,678	Lind et al. Manuscript
<u>Fathead minnow, <i>Pimephales promelas</i></u>	Nickel sulfate	91	96 hr	LC50 (TOC = 30 mg/L)	8,617	Lind et al. Manuscript
<u>Channel catfish ($1.2\text{-}1.5$ g), <i>Ictalurus punctatus</i></u>	Nickel sulfate	42	48 hr	LC50	36,840	Willford 1966
<u>Channel catfish, <i>Ictalurus punctatus</i></u>	Nickel chloride	93- 105	7 days	EC50 (death and deformity)	710	Birge and Black 1980; Birge et al. 1981

Table 6. (Continued)

<u>Species</u>	<u>Chemical</u>	<u>Hardness (mg/L as CaCO_3)</u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration ($\mu\text{g/L}$)^a</u>	<u>Reference</u>
Guppy, <i>Poecilia reticulata</i>	Nickel sulfate	260	96 hr	LC50 (high solids)	34,900	Khangarot 1981
Guppy (184 mg), <i>Poecilia reticulata</i>	Nickel chloride	260	48 hr	LC50	37,000	Khangarot et al. 1981
Bluegill (0.7-1.1 g), <i>Lepomis macrochirus</i>	Nickel sulfate	42	48 hr	LC50	110,500	Willford 1966
Largemouth bass (embryo, larva), <i>Micropterus salmoides</i>	Nickel chloride	93-105	8 days	EC50 (death and deformity)	2,020 (2,060)	Birge and Black 1980; Birge et al. 1978, 1981
Narrow-mouthed toad (embryo, larva), <i>Gastrophryne carolinensis</i>	Nickel chloride	195	7 days	EC50 (death and deformity)	50	Birge 1978; Birge et al. 1979
Narrow-mouthed toad (embryo, larva), <i>Gastrophryne carolinensis</i>	Nickel chloride	95-103	7 days	EC50 (death and deformity)	50	Birge and Black 1980
Fowler's toad, <i>Bufo fowleri</i>	Nickel chloride	93-105	7 days	EC50 (death and deformity)	11,030	Birge and Black 1980
Marbled salamander (embryo, larva), <i>Ambystoma opacum</i>	Nickel chloride	93-105	8 days	EC50 (death and deformity)	420 (410)	Birge and Black 1980; Birge et al. 1978

Table 6. (Continued)

<u>Species</u>	<u>Chemical</u>	<u>Salinity (g/kg)</u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration ($\mu\text{g/L}$)^a</u>	<u>Reference</u>
<u>SALTWATER SPECIES</u>						
Golden brown alga, <u><i>Isochrysis galbana</i></u>	-	28	2 days	Lowest concentration reducing chlorophyll a	500	Wilson and Freeberg 1980
Golden brown alga, <u><i>Isochrysis galbana</i></u>	-	28	9 days	Lowest concentration reducing cell numbers	80	Wilson and Freeberg 1980
Diatom, <u><i>Phaeodactylum tricornutum</i></u>	Nickel chloride	26	7 days	Reduced growth	1,000	Skaar et al. 1974
Diatom, <u><i>Thalassiosira pseudonana</i></u>	-	14	2 days	Chlorophyll a reduced about 65% at 12°C	100	Wilson and Freeberg 1980
Diatom, <u><i>Thalassiosira pseudonana</i></u>	-	14	2 days	Chlorophyll a reduced about 65% at 16°C	31	Wilson and Freeberg 1980
Diatom, <u><i>Thalassiosira pseudonana</i></u>	-	14	2 days	Chlorophyll a reduced about 65% at 20°C	28	Wilson and Freeberg 1980
Diatom, <u><i>Thalassiosira pseudonana</i></u>	-	14	2 days	Chlorophyll a reduced about 65% at 24°C	17	Wilson and Freeberg 1980
Diatom, <u><i>Thalassiosira pseudonana</i></u>	-	14	2 days	Chlorophyll a reduced about 65% at 28°C	80	Wilson and Freeberg 1980
Diatom, <u><i>Thalassiosira pseudonana</i></u>	-	28	2 days	Chlorophyll a reduced about 65% at 12°C	72	Wilson and Freeberg 1980
Diatom, <u><i>Thalassiosira pseudonana</i></u>	-	28	2 days	Chlorophyll a reduced about 65% at 16°C	140	Wilson and Freeberg 1980

Table 6. (Continued)

<u>Species</u>	<u>Chemical</u>	<u>Salinity (g/kg)</u>	<u>Duration</u>	<u>Effect</u>	<u>Concentration (µg/L)*</u>	<u>Reference</u>
<u>Sea urchin (gamete), Strongylocentrotus purpuratus</u>	-	-	300 min	Depressed sperm motility	58.69	Timour Ian and Watchmaker 1977

* Results are expressed as nickel, not as the chemical.

** Incubated for 2 to 4 days after exposure.

REFERENCES

Abbe, G.R. 1982. Growth, mortality and copper-nickel accumulation by oysters (Crassostrea virginica) at the Morgantown steam electric station on the Potomac River, Maryland. J. Shellfish Res. 2:3-13.

Abo-Rady, M. 1979. The levels of heavy metals (Cd, Cu, Hg, Ni, Pb, Zn) in brook trouts from the River Leine in the area of Gottingen (West Germany). Z. Lebensm. Unters. Forsch. 168:259-263.

Agrawal, S.J., A.K. Srivastava and H.S. Chaudhry. 1979. Haematological effects of nickel toxicity on a fresh water teleost, Colisa fasciatus. Acta Pharmacol. Toxicol. 45:215-217.

Ahsanullah, M. 1982. Acute toxicity of chromium, mercury, molybdenum, and nickel to the amphipod Allorchestes compressa. Aust. J. Mar. Freshwater Res. 33:465-474.

Albright, L.J., J.W. Wentworth and E.M. Wilson. 1972. Technique for measuring metallic salt effects upon the indigenous heterotrophic microflora of a natural water. Water Res. 6:1589-1596.

Alman, H. and P. Boger. 1984. Nickel-dependent uptake-hydrogenase activity in the blue-green alga Anabaena variabilis. Z. Naturforsch. 39C:90-92.

Amemiya, Y. and O. Nakayama. 1984. The chemical composition and metal absorption capacity of the sheath materials isolated from Microcystis, cyanobacteria. Jpn. J. Limnol. 45:187-193.

Anderson, B.G. 1948. The apparent thresholds of toxicity to Daphnia magna for chlorides of various metals when added to Lake Erie water. Trans. Am. Fish. Soc. 78:96-113.

Anderson, D.R. 1981. The combined effects of nickel, chlorine, and temperature on the mortality of rainbow trout, Salmo gairdneri. Ph.D. thesis. University of Washington, Seattle, WA. Available from: University Microfilms, Ann Arbor, MI. Order No. 8121178.

Anderson, D.R. 1983. Chlorine-heavy metals interaction on toxicity and metal accumulation. In: Water chlorination: Environmental impact and health effects. Vol. 4, Book 2. Jolley, R.L., W.A. Brungs, J.A. Cotruvo, R.B. Cumming, J.S. Mattice and V.A. Jacobs (Eds.). Ann Arbor Science Publishers, Ann Arbor, MI. pp. 811-826.

Anderson, P.D. 1973. An approach to the study of multiple toxicity through the derivation and use of quantal response curves. Ph.D. thesis. Oregon State University, Corvallis, OR. Available from: University Microfilms, Ann Arbor, MI. Order No. 74-4187.

Anderson, P.D. and L.J. Weber. 1975. Toxic response as a quantitative function of body size. Toxicol. Appl. Pharmacol. 33:471-483.

Applegate, V.C., J.H. Howell, A.E. Hall, Jr. and M.A. Smith. 1957. Toxicity of 4,346 chemicals to larval lampreys and fishes. Special Scientific Report-- Fisheries No. 207. U.S. Fish and Wildlife Service, Washington, DC.

Arillo, A., C. Margiocco, F. Melodia and P. Mensi. 1982. Biochemical effects of long term exposure to Cr, Cd, Ni on rainbow trout (Salmo gairdneri Rich.): Influence of sex and season. Chemosphere 11:47-57.

Babich, H. and G. Stotzky. 1983. Temperature, pH, salinity, hardness, and particulates mediate nickel toxicity to eubacteria, an actinomycete, and yeasts in lake, simulated estuarine, and sea waters. Aquat. Toxicol. 3:195-208.

Babich, H. and G. Stotzky. 1985. Heavy metal toxicity to microbe-mediated ecologic processes: A review and potential application to regulatory policies. Environ. Res. 36:111-137.

Babich, H., C. Shopsis and E. Borenfreund. 1986. In vitro cytotoxicity testing of aquatic pollutants (cadmium, copper, zinc, nickel) using established fish cell lines. Ecotoxicol. Environ. Safety 11:91-99.

Bailey, R.C. and P.M. Stokes. 1985. Evaluation of filamentous algae as biomonitor of metal accumulation in softwater lakes: A multivariate approach. In: Aquatic toxicology and hazard assessment: 7th symposium. Cardwell, R.D., R. Purdy and R.C. Bahner (Eds.). ASTM STP 854. American Society for Testing and Materials, Philadelphia, PA. pp. 5-26.

Ballester, A. and J. Castellvi. 1979. Contribution to the biokinetic study of vanadium and nickel uptake by marine organisms. Invest. Pesq. 43:449-478.

Baudouin, M.F. and P. Scoppa. 1974. Acute toxicity of various metals to freshwater zooplankton. Bull. Environ. Contam. Toxicol. 12:745-751.

Bean, B. and A. Harris. 1977. A calcium-reversible selective inhibition of flagellar coordination by nickel ion in Chlamydomonas. Abstracts of the annual meeting of the American Society of Microbiology.

Becker, C.D. and M.G. Wolford. 1980. Thermal resistance of juvenile salmonids sublethally exposed to nickel, determined by the critical thermal maximum method. Environ. Pollut. (Series A) 21:181-189.

Bengtsson, B. 1978. Use of a harpacticoid copepod in toxicity tests. Mar. Pollut. Bull. 9:238-241.

Besser, J.M. 1985. Bioavailability and toxicity of heavy metals in mine tailings leachate to aquatic invertebrates. M.S. thesis. University of Missouri, Columbia, MS.

Biddinger, G.R. and S.P. Gloss. 1984. The importance of trophic transfer in bioaccumulation of chemical contaminants in aquatic ecosystems. Residue Rev. 91:103-145.

Biesinger, K.E. and G.M. Christensen. 1972. Effect of various metals on survival, growth, reproduction, and metabolism of Daphnia magna. J. Fish. Res. Board Can. 29:1691-1700.

Birge, W.J. 1978. Aquatic toxicology of trace elements and coal and fly ash. In: Energy and environmental stress in aquatic systems. Thorp, J.H. and J.W. Gibbons (Eds.). CONF-771114. National Technical Information Service, Springfield, VA. pp. 219-240.

Birge, W.J. and J.A. Black. 1980. Aquatic toxicology of nickel. In: Nickel in the environment. Nriagu, J.O. (Ed.). Wiley, New York, NY. pp. 349-366.

Birge, W.J., J.E. Hudson, J.A. Black and A.G. Westerman. 1978. Embryo-larval bioassays on inorganic coal elements and in situ biomonitoring of coal-waste effluents. In: Surface mining and fish/wildlife needs in the eastern United States. Samuel, D.E., J.R. Stauffer, C.H. Hocutt and W.T. Mason (Eds.). PB298353. National Technical Information Service, Springfield, VA. pp. 97-104.

Birge, W.J., J.A. Black and A.G. Westerman. 1979. Evaluation of aquatic pollutants using fish and amphibian eggs as bioassay organisms. In: Animals as monitors of environmental pollutants. Neilson, S.W., G. Migaki and D.G. Scarrelli (Eds.). National Academy of Sciences, Washington, DC. pp. 108-118.

Birge, W.J., J.A. Black, A.G. Westerman and J.E. Hudson. 1980. Aquatic toxicity tests on inorganic elements occurring in oil shale. In: Oil shale symposium: Sampling, analysis and quality assurance. Gale, C. (Ed.). PB80-221435 or EPA-600/9-80-022. National Technical Information Service, Springfield, VA. pp. 519-534.

Birge, W.J., J.A. Black, and B.A. Ramey. 1981. The reproductive toxicology of aquatic contaminants. In: Hazard assessment of chemicals: Current developments. Vol. 1. Saxena, J. and F. Fisher (Eds.). Academic Press, New York, NY. pp. 59-115.

Blaise, C. and P. Couture. 1984. Detection with the aid of bioassay (Selenastrum capricornutum) of the effects of trace metals in the aquatic environment. Toxicity or enrichment by essential nutrients. Hydrobiologia 114:39-50.

Blaylock, B.G. and M.L. Frank. 1979. A comparison of the toxicity of nickel to the developing eggs and larvae of carp (Cyprinus carpio). Bull. Environ. Contam. Toxicol. 21:604-611.

Bornatowicz, N. 1983. Assessment of the acute toxicity of nickel sulfate to rainbow trout. PB84232073. National Technical Information Service, Springfield, VA.

Bosserman, R.W. 1985. Distribution of heavy metals in aquatic macrophytes from Okefenokee swamp. Symp. Biol. Hung. 29:31-40.

Bradley, R.W. and J.R. Morris. 1986. Heavy metals in fish from a series of metal-contaminated lakes near Sudbury, Ontario. Water Air Soil Pollut. 27:341-354.

Braginskiy, L.P. and E.P. Shcherban. 1978. Acute toxicity of heavy metals to aquatic invertebrates at different temperatures. *Hydrobiol. J.* 14(6):78-82.

Brezina, E.R. and M.Z. Arnold. 1977. Levels of heavy metals in fishes from selected Pennsylvania waters. Publication No. 50. Pennsylvania Department of Environmental Resources, Bureau of Water Quality Management, Harrisburg, PA.

Bringmann, G. 1978. Determination of the biological toxicity of waterbound substances towards protozoa. I. Bacteriovorous flagellates (model organism: Entosiphon sulcatum Stein). *Z. Wasser Abwasser Forsch.* 11:210-215.

Bringmann, G. and R. Kuhn. 1959a. The toxic effects of waste water on aquatic bacteria, algae, and small crustaceans. *Gesund.-Ing.* 80:115-120.

Bringmann, G. and R. Kuhn. 1959b. Water toxicology studies with protozoans as test organisms. *Gesund.-Ing.* 80:239-242.

Bringmann, G. and R. Kuhn. 1977a. Limiting values for the damaging action of water pollutants to bacteria (Pseudomonas putida) and green algae (Scenedesmus quadricauda) in the cell multiplication inhibition test. *Z. Wasser Abwasser Forsch.* 10:87-98.

Bringmann, G. and R. Kuhn. 1977b. Results of the damaging effect of water pollutants on Daphnia magna. *Z. Wasser Abwasser Forsch.* 10:161-166.

Bringmann, G. and R. Kuhn. 1978a. Limiting values for the noxious effects of water pollutant material to blue algae (*Microcystis aeruginosa*) and green algae (*Scenedesmus quadricauda*) in cell propagation inhibition tests. Vom Wasser 50:45-60.

Bringmann, G. and R. Kuhn. 1978b. Testing of substances for their toxicity threshold: Model organisms *Microcystis (Diplocystis) aeruginosa* and *Scenedesmus quadricauda*. Mitt. Int. Ver. Theor. Angew. Limnol. 21:275-284.

Bringmann, G. and R. Kuhn. 1979. Comparison of toxic limiting concentrations of water contamination toward bacteria, algae and protozoa in the cell-growth inhibition test. Haustech. Bauphys. Umwelttech. 100:249-252.

Bringmann, G. and R. Kuhn. 1980a. Determination of the harmful biological effect of water pollutants on protozoa. II. Bacteriovorous ciliates. Z. Wasser Abwasser Forsch. 13:26-31.

Bringmann, G. and R. Kuhn. 1980b. Comparison of the toxicity thresholds of water pollutants to bacteria, algae, and protozoa in the cell multiplication inhibition test. Water Res. 14:231-241.

Bringmann, G. and R. Kuhn. 1981. Comparison of the effects of harmful substances on flagellates as well as ciliates and on halozoic bacteriophagous and saprozoic protozoa. Gas-Wasserfach, Wasser-Abwasser 122:308-313.

Bringmann, G. and R. Kuhn. 1982. Results of toxic action of water pollutants on Daphnia magna Straus tested by an improved standardized procedure. Z. Wasser Abwasser Forsch 15:1-6.

Bringmann, G., R. Kuhn and A. Winter. 1980. Determination of biological damage from water pollutants to protozoa. III. Saprozoic flagellates. Z. Waasser Abwasser Forsch. 13:170-173.

Brkovic-Popovic, I. and M. Popovic. 1977a. Effects of heavy metals on survival and respiration rate of tubificid worms: Part I - Effects on survival. Environ. Pollut. 13:65-72.

Brkovic-Popovic, I. and M. Popovic. 1977b. Effects of heavy metals on survival and respiration rate of tubificid worms: Part II - Effects on respiration rate. Environ. Pollut. 13:93-98.

Brown, B.T. and B.M. Rattigan. 1979. Toxicity of soluble copper and other metal ions to Elodea canadensis. Environ. Pollut. 20:303-314.

Brown, V.M. 1968. The calculation of the acute toxicity of mixtures of poisons to rainbow trout. Water Res. 2:723-733.

Brown, V.M. and R.A. Dalton. 1970. The acute lethal toxicity to rainbow trout of mixtures of copper, phenol, zinc and nickel. J. Fish Biol. 2:211-216.

Bryan, G.W., W.J. Langston, L.G. Hummerstone, G.R. Burt and Y.B. Ho. 1983. An assessment of the gastropod, Littorina littorea, as an indicator of heavy metal contamination in United Kingdom estuaries. J. Mar. Biol. Assoc. U.K. 63:327-345.

Bryant, V., O.M. Newberry, A.S. McLusky and R. Campbell. 1985. Effect of temperature and salinity on the toxicity of nickel and zinc to two estuarine invertebrates (Corophium volutator, Macoma balthica). Mar. Ecol. Prog. Ser. 24:139-153.

Buikema, A.L., Jr., J. Cairns, Jr. and G.W. Sullivan. 1974a. Rotifers as monitors of heavy metal pollution in water. Bulletin 71. Virginia Water Resources Research Center, Blacksburg, VA.

Buikema, A.L., Jr., J. Cairns, Jr. and G.W. Sullivan. 1974b. Evaluation of Philodina acuticornis (Rotifera) as a bioassay organism for heavy metals. Water Resour. Bull. 10:648-661.

Buikema, A.L., Jr., C.L. See and J. Cairns, Jr. 1977. Rotifer sensitivity to combinations of inorganic water pollutants. Bulletin 92. Virginia Water Resources Research Center, Blacksburg, VA.

Cairns, J., Jr., K.W. Thompson and A.C. Hendricks. 1981. Effects of fluctuating, sublethal applications of heavy metal solutions upon the gill ventilation response of bluegills (Lepomis macrochirus). EPA-600/3-81-003. National Technical Information Service, Springfield, VA.

Calabrese, A. and D.A. Nelson. 1974. Inhibition of embryonic development of the hard clam, Mercenaria mercenaria by heavy metals. Bull. Environ. Contam. Toxicol. 11:92-97.

Calabrese, A., R.S. Collier, D.A. Nelson and J.R. MacInnes. 1973. The toxicity of heavy metals to embryos of the American oyster Crassostrea virginica. Mar. Biol. (Berl.) 18:162-166.

Calabrese, A., J.R. MacInnes, D.A. Nelson and J.E. Miller. 1977. Survival and growth of bivalve larvae under heavy-metal stress. Mar. Biol. (Berl.) 41:179-184.

Calamari, D., G.F. Gaggino and G. Pacchetti. 1982. Toxicokinetics of low levels of Cd, Cr, Ni and their mixture in long-term treatment on Salmo gairdneri Rich. Chemosphere 11:59-70.

Call, D.J., L.T. Brooke, N. Ahmad and J.E. Richter. 1983. Toxicity and metabolism studies with EPA priority pollutants and related chemicals in freshwater organisms. PB83263665 or EPA-600/3-83-095. National Technical Information Service, Springfield, VA.

Callahan, M.A., M.W. Slimak, N.W. Gabel, I.P. May, C.F. Fowler, J.R. Freed, P. Jennings, R.L. Durfee, F.C. Whitmore, B. Maestri, W.R. Mabey, B.R. Holt and C. Gould. 1979. Water-related environmental fate of 129 priority pollutants. Vol I. EPA-440/4-79-029a. National Technical Information Service, Springfield, VA. pp. 15-1 to 15-9.

Campbell, P.M. and G.D. Smith. 1986. Transport and accumulation of nickel ions in the cyanobacterium Anabaena cylindrica. Arch. Biochem. Biophys. 244:470-477.

Cardin, J.A. 1985. U.S. EPA, Narragansett, RI. (Memorandum to D.J. Hansen. U.S. EPA, Narragansett, RI.)

Chapman, G.A., S. Ota and F. Recht. Manuscript. Effects of water hardness on the toxicity of metals to Daphnia magna. U.S. EPA, Corvallis, OR.

Chapman, P.M. 1985. Effects of gut sediment contents on measurements of metal levels in benthic invertebrates - a cautionary note. Bull. Environ. Contam. Toxicol. 35:345-347.

Chapman, W.H., H.L. Fisher and M.W. Pratt. 1968. Concentration factors of chemical elements in edible aquatic organisms. UCRL-50564. National Technical Information Service, Springfield, VA.

Chassard-Bouchard, C. and G. Balvay. 1978. Application of electron probe x-ray microanalysis to the detection of metal pollutants in freshwater zooplankton. Microsc. Acta (Suppl.) 2:185-192.

Chaudhry, H.S. 1984. Nickel toxicity on carbohydrate metabolism of a freshwater fish, Colisa fasciatus. *Toxicol. Lett.* 20:115-121.

Chaudhry, H.S. and K. Nath. 1985. Nickel induced hyperglycemia in the freshwater fish, Colisa fasciatus. *Water Air Soil Pollut.* 24:173-176.

Chaudry, H.S. and K. Nath. 1985. Effect of nickel intoxication on liver glycogen of a freshwater teleost, Colisa fasciatus. *Acta Hydrochim. Hydrobiol.* 13:245-248.

Cherry, D.S., R.K. Guthrie, F.F. Sherberger and S.R. Lerrick. 1979. The influence of coal ash and thermal discharges upon the distribution and bioaccumulation of aquatic invertebrates. *Hydrobiologia* 62:257-267.

Christensen, E.R., C.Y. Chen and J. Kannall. 1985. The response of aquatic organisms to mixtures of toxicants. *Water Sci. Technol.* 17:1445-1446.

Clendenning, K.A. and W.J. North. 1959. Effects of wastes on the giant kelp, Macrocystis pyrifera. In: *Effects of wastes on the giant kelp, Macrocystis pyrifera*. Pearson, E.A. (Ed.). Proceedings of the first international conference on waste disposal in the marine environment, Berkeley, CA.

Cowgill, U.M., H.W. Emmel and I.T. Takahashi. 1985. Inorganic chemical composition of trout food pellets and alfalfa used to sustain Daphnia magna Straus. *Bull. Environ. Contam. Toxicol.* 34:890-896.

Cowgill, U.M., H.W. Emmel, D.L. Hopkins, S.L. Applegath and I.T. Takahashi. 1986. The influence of water on reproductive success and chemical composition of laboratory reared populations of Daphnia magna. Water Res. 20:317-323.

Daday, A., A.H. Mackerras and G.D. Smith. 1985. The effect of nickel on hydrogen metabolism and nitrogen fixation in the cyanobacterium Anabaena cylindrica. J. Gen. Microbiol. 131:231-238.

Dallinger, R. and H. Kautzky. 1985. The importance of contaminated food for the uptake of heavy metals by rainbow trout (Salmo gairdneri): A field study. Oecologia 67:82-89.

Devi Prasad, P.V. and P.S. Devi Prasad. 1982. Effect of cadmium, lead and nickel on three freshwater green algae. Water Air Soil Pollut. 17:263-268.

Dixon, W.J. and M.B. Brown (Eds.). 1979. BMDP Biomedical Computer Programs, P-series. University of California, Berkeley, CA. p. 521.

Dorfman, D. 1977. Tolerance of Fundulus heteroclitus to different metals in salt water. Bull. N.J. Acad. Sci. 22:21-23.

Doudoroff, P. 1956. Some experiments on the toxicity of complex cyanides to fish. Sewage Ind. Wastes. 28:1020-1026.

Doudoroff, P. and M. Katz. 1953. Critical review of literature on the toxicity of industrial wastes and their components to fish. II. The metals, as salts. 25:802-839.

Doudoroff, P., G. Leduc and C.R. Schneider. 1966. Acute toxicity to fish of solutions containing complex metal cyanides, in relation to concentrations of molecular hydrocyanic acid. Trans. Am. Fish. Soc. 95:6-22.

Dugan, P.R. 1975. Bioflocculation and the accumulation of chemicals by floc-forming organisms. EPA 600/2-75-032. National Technical Information Service, Springfield, VA.

Dunstan, J.C., A. deForest and R.W. Pettis. 1980. Mytilus edulis as an indicator of trace metal pollution in naval dockyard waters with preliminary results from Williamstown Naval Dockyards, Victoria. MRL-R-781. Department of Defense, Materials Research Laboratories, Melbourne, Victoria, Canada.

Eisler, R. 1977a. Acute toxicities of selected heavy metals to the soft-shell clam, Mya arenaria. Bull. Environ. Contam. Toxicol. 17:137-145.

Eisler, R. 1977b. Toxicity evaluation of a complex metal mixture to the softshell clam Mya arenaria. Mar. Biol. (Berl.) 43:265-276.

Eisler, R. 1981. Trace metal concentrations in marine organisms. Pergamon Press, New York, NY.

Eisler, R. and R.J. Hennekey. 1977. Acute toxicities of Cd^{+2} , Cr^{+6} , Hg^{+2} , Ni^{+2} and Zn^{+2} to estuarine macrofauna. Arch. Environ. Contam. Toxicol. 6:315-323.

Eisler, R., M.M. Barry, R.I. Lapan, Jr., G. Telek, E.W. Davey and A.E. Soper. 1978. Metal survey of the marine clam Pifid' Botrhina collected near a Rhode Island (USA) electroplating plant. Mar. Biol. (Berl.) 45:311-317.

Ellis, M.M. 1937. Detection and measurement of stream-pollution. Bulletin No. 22. Bureau of Fisheries, U.S. Department of Commerce, Washington, DC.

Fezy, J.S., D.F. Spencer and R.W. Greene. 1979. The effect of nickel on the growth of the freshwater diatom Navicula pelliculosa. Environ. Pollut. 20:131-137.

Forstner, U. 1984. Metal pollution of terrestrial waters. In: Changing metal cycles and human health. Nriagu, J.O. (Ed.). Springer-Verlag, New York, NY. pp. 71-94.

Foster, P. 1976. Concentration and concentration factors of heavy metals in brown algae. Environ. Pollut. 10:9-17.

Gentile, J.H., S.M. Gentile, N.G. Hairston, Jr. and B.K. Sullivan. 1982. The use of life-tables for evaluating the chronic toxicity of pollutants to Mysidopsis bahia. Hydrobiologia 93:179-187.

Gerhards, U. and H. Weller. 1977. The uptake of mercury, cadmium and nickel by Chlorella pyrenoidosa. Z. Pflanzenphysiol. 82:292-300.

Giattina, J.D., R.R. Garton and D.G. Stevens. 1982. Avoidance of copper and nickel by rainbow trout as monitored by a computer-based data acquisition system. Trans. Am. Fish. Soc. 111:491-504.

Gill, T.S. and J.C. Pant. 1981. Toxicity of nickel to the fish Puntius conchonius (Ham.) and its effects on blood glucose and liver glycogen. Comp. Physiol. Ecol. 6:99-102.

Gordon, M., G.A. Knauer and J.H. Martin. 1980. Mytilus californianus as a bioindicator of trace metal pollution: Variability and statistical considerations. Mar. Pollut. Bull. 11:195-198.

Grande, M. and S. Andersen. 1983. Lethal effects of hexavalent chromium, lead and nickel on young stages of Atlantic salmon (Salmo salar L.) in soft water. Vatten 39:405-416.

Guilizzoni, P. 1980. Heavy metals distribution and their effects on photosynthetic rates of two submerged macrophytes of Lake Mezzola. In: Environment and quality of life. Second environmental research programme 1976-1980. EUR-6388 EN. ISBN 92-825-0496-4. Commission of the European Communities, Brussels, Luxembourg. pp. 34-38.

Hale, J.G. 1977. Toxicity of metal mining wastes. Bull. Environ. Contam. Toxicol. 17:66-73.

Hall, R.A., E.G. Zook and G.M. Meaburn. 1978. National Marine Fisheries Service survey of trace elements in the fishery resource. NOAA Technical Report NMFS SSRF-721. National Technical Information Service, Springfield, VA.

Hall, T. 1978. Nickel uptake, retention and loss in Daphnia magna. M.S. thesis. University of Toronto, Toronto, Ontario, Canada.

Hall, T.M. 1982. Free ionic nickel accumulation and localization in the freshwater zooplankter, Daphnia magna. Limnol. Oceanogr. 27:718-727.

Hansen, D.J. 1983. U.S. EPA, Gulf Breeze, FL. (Memorandum to W.A. Brungs. U.S. EPA, Narragansett, RI.)

Hardy, J.T. and G. Roesijadi. 1982. Bioaccumulation kinetics and oxygen distribution of nickel in the marine clam Prototthaca staminea. Bull. Environ. Contam. Toxicol. 28:566-572.

Havas, M. and T.C. Hutchinson. 1982. Aquatic invertebrates from the Smoking Hills, N.W.T.: Effect of pH and metals on mortality. Can. J. Fish. Aquat. Sci. 39:890-903.

Heit, M. and C.S. Klusek. 1985. Trace element concentrations in the dorsal muscle of white suckers and brown bullheads from two acidic Adirondack lakes. Water Air Soil Pollut. 25:87-96.

Ho, M.S. and P.L. Zubkoff. 1983. The opposing effects of cadmium and nickel on calcium uptake by larvae of the clam Mulinia lateralis. Comp. Biochem. Physiol. 74C:337-339.

Hughes, G.M., S.F. Perry and V.M. Brown. 1979. A morphometric study of effects of nickel, chromium and cadmium on the secondary lamellae of rainbow trout gills. *Water Res.* 13:665-679.

Hutchinson, N.J. and J.B. Sprague. 1983. Chronic toxicity of a mixture of 7 metals to flagfish in soft, acid water. In: *Proceedings of the eighth annual aquatic toxicity workshop*. Kaushik, N.K. and K.R. Solomon (Eds.). Canadian Technical Report of Fisheries and Aquatic Sciences No. 1151. *Fisheries and Oceans*, Ottawa, Ontario, Canada. p. 191.

Hutchinson, J.N. and J.B. Sprague. 1986. Toxicity of trace metal mixtures to American flagfish (*Jordanella floridae*) in soft, acidic water and implications for cultural acidification. *Can. J. Fish. Aquat. Sci.* 43:647-655.

Hutchinson, T.C. 1973. Comparative studies of the toxicity of heavy metals to phytoplankton and their synergistic interactions. *Water Pollut. Res. Can.* 8:68-90.

Hutchinson, T.C. and P.M. Stokes. 1975. Heavy metal toxicity and algal bioassays. In: *Water quality parameters*. Barabas, S. (Ed.). ASTM STP 573. American Society for Testing and Materials, Philadelphia, PA. pp. 320-343.

Hutchinson, T.C., A. Fedorenko, J. Fitchko, A. Kuja, J. Van Loon and J. Lichwa. 1975. Movement and compartmentation of nickel and copper in an aquatic ecosystem. In: *Trace substances in environmental health - IX*. Hemphill, D.D. (Ed.). University of Missouri, Columbia, MO. pp. 89-105.

Janauer, G.A. 1985. Heavy metal accumulation and physiological effects on Austrian macrophytes. *Symp. Biol. Hung.* 29:21-30.

Jay, F.B. and R.J. Muncy. 1979. Toxicity to channel catfish of wastewater from an Iowa coal beneficiation plant. *Iowa State J. Res.* 54:45-50.

Jenkins, D.W. 1980. Biological monitoring of toxic trace metals, Vol. 2: Toxic trace metals in plants and animals of the world. Part III. EPA-600/3-80-092 or PB81-103509. National Technical Information Service, Springfield, VA.

Jennett, J.C., J.E. Smith and J.M. Hassett. 1982. Factors influencing metal accumulation by algae. EPA-600/2-82-100. National Technical Information Service, Springfield, VA.

Jones, J.R.E. 1935. The toxic action of heavy metal salts on the three-spined stickleback (Gasterosteus aculeatus). *J. Exp. Biol.* 12:165-173.

Jones, J.R.E. 1939. The relation between the electrolytic solution pressures of the metals and their toxicity to the stickleback (Gasterosteus aculeatus L.). *J. Exp. Biol.* 16:425-437.

Kaiser, K.L.E. 1980. Correlation and prediction of metal toxicity to aquatic biota. *Can. J. Fish. Aquat. Sci.* 37:211-218.

Kanai, K. and H. Wakabayashi. 1984. Purification and some properties of protease from Aeromonas hydrophila. Bull. Jpn. Soc. Sci. Fish. 50:1367-1374.

Kapur, K. and N.A. Yadav. 1982. The effects of certain heavy metal salts on the development of eggs in common carp, Cyprinus carpio var. communis. Acta Hydrochim. Hydrobiol. 10:517-522.

Kawamata, S., Y. Yamauro, H. Hayashi and S. Komiyama. 1983. Contents of heavy metals in fishes in Nagano prefecture. CA Selects: Environ. Pollut. Issue 7:4. Abstr. No. 98:105850d, 1983.

Keller, W. and J.R. Pitblado. 1984. Crustacean plankton in northeastern Ontario lakes subjected to acidic deposition. Water Air Soil Pollut. 23:27-291.

Khangarot, B.S. 1981. Lethal effects of zinc and nickel on freshwater teleosts. Acta Hydrochim. Hydrobiol. 9:297-302.

Khangarot, B.S., V.S. Durve and V.K. Rajbanshi. 1981. Toxicity of interactions of zinc-nickel, copper-nickel and zinc-nickel-copper to a freshwater teleost, Lebiasina reticulatus (Peters). Acta Hydrochim. Hydrobiol. 9:495-503.

Khangarot, B.S., S. Mathur, and V.S. Durve. 1982. Comparative toxicity of heavy metals and interaction of metals on a freshwater pulonate snail Lymnaea acuminata (Lamarck). Acta Hydrochim. Hydrobiol. 10:367-375.

Kissa, E., M. Moraitou-Apostolopoulou and V. Kiortsis. 1984. Effects of four heavy metals on survival and hatching of Artemia salina (L.). Arch. Hydrobiol. 102:255-264.

Kopp, J.F. and R.C. Kroner. 1967. Trace metals in waters of the United States, Oct. 1, 1962, to Sept. 30, 1967. Federal Water Pollution Administration, Cincinnati, OH.

Kumar, D., M. Jha and H.D. Kumar. 1985. Heavy-metal toxicity in the cyanobacterium Nostoc linckia. Aquat. Bot. 22:101-105.

La Touche, Y.D. and M.C. Mix. 1982. Seasonal variations of arsenic and other trace elements in bay mussels (Mytilus edulis). Bull. Environ. Contam. Toxicol. 29:665-670.

Lazareva, L.P. 1985. Changes in biological characteristics of Daphnia magna from chronic action of copper and nickel at low concentrations. Hydrobiol. J. 21(5):59-62.

LeBlanc, G.A. 1984. Interspecies relationships in acute toxicity of chemicals to aquatic organisms. Environ. Toxicol. Chem. 3:47-60.

Lewis, M.A. 1986. Impact of municipal wastewater effluent on water quality, periphyton, and invertebrates in the Little Miami River near Xenia, Ohio. Ohio J. Sci. 86:2-8.

Lind, D., K. Alto and S. Challderston. Manuscript. Regional copper-nickel study: Aquatic toxicological study. Minnesota Environmental Quality Board, Minneapolis, MN.

Lopez-Avila, V., W.D. McKenzie, W.W. Sutton, R. Kaminsky, U. Spanagel, T.A. Olsson and J.H. Taylor. 1985. Application of chemical fractionation/aquatic bioassay procedure to hazardous waste site monitoring. EPA-600/4-85-059 or PB86-109493. National Technical Information Service, Springfield, VA.

Lorz, H.W., R.H. Williams and C.A. Fustish. 1978. Effects of several metals on smolting of coho salmon. EPA-600/3-78-090. National Technical Information Service, Springfield, VA.

Lussier, S.M. and J.A. Cardin. 1985. U.S. EPA, Narragansett, RI. (Memorandum to D.J. Hansen. U.S. EPA, Narragansett, RI.)

Lussier, S.M., J.H. Gentile and J. Walker. 1985. Acute and chronic effects of heavy metals and cyanide on Mysidopsis bahia (Crustacea: Mysidacea). Aquat. Toxicol. 7:25-35.

MacDonald, C.R., T.R. Jack and J.E. Zajic. 1980. Metal ion toxicity to Daphnia magna: A novel approach to data plotting. In: Proceedings of the sixth annual aquatic toxicity workshop. Klaverkamp, J.F., S.L. Leonhard and K.E. Marshall (Eds.). Canadian technical report of fisheries and aquatic sciences No. 975. Department of Fisheries and Oceans, Ottawa, Ontario, Canada. pp. 91-107.

Malueg, K.W., G.S. Schuytema, J.H. Gakstatter and D.F. Krawczyk. 1984.
Toxicity of sediments from three metal-contaminated areas. Environ. Toxicol. Chem. 3:279-291.

Mann, H. and W.S. Fyfe. 1984. An experimental study of algal uptake of U, Ba, V, Co and Ni from dilute solutions. Chem. Geol. 44:385-398.

Mansouri-Aliabadi, M. and R.E. Sharp. 1985. Passage of selected heavy metals from Sphaerotilus (Bacteria: Chlamydobacteriales) to Paramecium caudatum (Protozoa: Ciliasta). Water Res. 19:697-699.

Markarian, R.K., M.C. Matthews and L.I. Connor. 1980. Toxicity of nickel, copper, zinc and aluminum mixtures to the white sucker (Catostomus commersoni). Bull. Environ. Contam. Toxicol. 25:790-796.

Martin, J.H. 1979. Bioaccumulation of heavy metals by littoral and pelagic marine organisms. EPA-600/3-79-038. National Technical Information Service, Springfield, VA.

Martin, J.H. and G.A. Knauer. 1972. A comparison of inshore versus offshore levels of twenty-one trace and major elements in marine plankton. In: Baseline studies of pollutants in the marine environment (heavy metals, halogenated hydrocarbons and petroleum). Goldberg, E.D. (Ed.). Brookhaven National Laboratory. pp. 35-66.

Mathis, B.J. and T.F. Cummings. 1973. Selected metals in sediments, water, and biota in the Illinois River. J. Water Pollut. Control Fed. 45:1573-1583.

McCabe, L.J., J.M. Symons, R.D. Lee and G.G. Robeck. 1970. Survey of community water supply systems. J. Am. Water Works Assoc. 62:670-687.

McDermott, D.J., G.V. Alexander, D.R. Young and A.T. Mearns. 1976. Metal contamination of flatfish around a large submarine outfall. J. Water Pollut. Control Fed. 48:1913-1918.

McFeters, G.A., P.J. Bond, S.B. Olson and Y.T. Tchan. 1983. A comparison of microbial bioassays for the detection of aquatic toxicants. Water Res. 17:1757-1762.

McHardy, B.M. and J.J. George. 1985. The uptake of selected heavy metals by the green alga Cladophora glomerata. Symp. Biol. Hung. 29:2-19.

McKim, J.M. 1977. Evaluation of tests with early life stages of fish for predicting long-term toxicity. J. Fish. Res. Board Can. 34:1148-1154.

Mears, H.C. and R. Eisler. 1977. Trace metals in liver from bluefish, tautog and filefish in relation to body length. Chesapeake Sci. 18:315-318.

Michnowicz, C.J. and T.E. Weakley. 1984. Effects of pH on toxicity of As, Cr, Cu, Ni and Zn to Selenastrum capricornutum Printz. Hydrobiologia 118:299-305.

Muramoto, S. 1983. Influence of complexans (NTA, EDTA) on the toxicity of nickel chloride and sulfate to fish at high concentrations. J. Environ. Sci. Health 18A:787-795.

Muramoto, S. and Y. Oki. 1984. Influence of anionic surface-active agents on the uptake of heavy metals by water hyacinth (Eichhornia crassipes). Bull. Environ. Contam. Toxicol. 33:444-450.

Mushak, P. 1980. Metabolism and systemic toxicity of nickel. In: Nickel in the environment. Nriagu, J.O. (Ed.). Wiley, New York, NY. pp. 499-523.

Muska, C.F. 1978. Evaluation of an approach for studying the quantitative response of whole organisms to mixtures of environmental toxicants. Ph.D. thesis. Oregon State University, Corvallis, OR. Available from: University Microfilms, Ann Arbor, MI. Order No. 77-29,421.

Muska, C.F. and L.J. Weber. 1977a. An approach for studying the effects of mixtures of environmental toxicants on whole organism performances. In: Recent advances in fish toxicology. Tubb, R.A. (Ed.). PB273500 or EPA-600/3-77-085. National Technical Information Service, Springfield, VA.

Muska, C.F. and L.J. Weber. 1977b. An approach for studying the effects of mixtures of toxicants. Proc. West. Pharmacol. Soc. 20:427-430.

Nebeker, A.V., C. Savonen, R.J. Baker and J.K. McCrady. 1984. Effects of copper, nickel, and zinc on the life cycle of the caddisfly Clistoronia magnifica (Limnephilidae). Environ. Toxicol. Chem. 3:645-649.

Nebeker, A.V., C. Savonen and D.G. Stevens. 1985. Sensitivity of rainbow trout early life stages to nickel chloride. Environ. Toxicol. Chem. 4:233-239.

Neter, J. and W. Wasserman. 1974. Applied linear statistical models. Irwin, Inc., Homewood, IL.

Nriagu, J.O. 1980. Cilohal cycle and properties of nickel. In: Nickel in the environment. Nriagu, J.O. (Ed.). Wiley, New York, NY. pp. 1-26.

O'Connor, T.P. 1976. Investigation of heavy metal concentrations of sediment and biota in the vicinity of the Morgantown steam electric generating station. Morgantown Monitoring Program Report Series.

Ozimek, T. 1985. Heavy metal content in macrophytes from ponds supplied with post-sewage water. Symp. Biol. Hung. 29:41-50.

Palawski, D., J.B. Hunn and F.J. Dwyer. 1985. Sensitivity of young striped bass to organic and inorganic contaminants in fresh and saline waters. Trans. Am. Fish. Soc. 114:748-753.

Parsons, T.R., C.A. Bowden and W.A. Heath. 1972. Preliminary survey of mercury and other metals contained in animals from the Fraser River mudflats. J. Fish. Res. Board Can. 30:1014-1016.

Patrick, R., T. Bott and R. Larson. 1975. The role of trace elements in management of nuisance growths. PB241985 or EPA-660/2-75-008. National Technical Information Service, Springfield, VA.

Pennington, C.H., J.A. Baker and M.E. Potter. 1982. Contaminant levels in fishes from Brown's Lake, Mississippi. J. Miss. Acad. Sci. 27:139-147.

Petrich, S.M. and D.J. Reish. 1979. Effects of aluminum and nickel on survival and reproduction in polychaetous annelids. Bull. Environ. Contam. Toxicol. 23:698-702.

Petukhov, S.A. and E.M. Ninonenko. 1982. On the priority of toxicological "hazard" of nickel in the sea. Mar. Pollut. Bull. 12:426.

Phelps, D.K., W. Galloway, F.P. Thurberg, E. Gould and M.A. Dawson. 1981. Comparison of several physiological monitoring techniques as applied to the blue mussel, Mytilus edulis, along a gradient of pollutant stress in Narragansett Bay, Rhode Island. In: Biological monitoring of marine pollutants. Vernberg, F.J., A. Calabrese, F.P. Thurberg and W.B. Vernberg (Eds.). Academic Press, New York, NY. pp. 335-355.

Phillips, G.R. and R.C. Russo. 1978. Metal bioaccumulation in fishes and aquatic invertebrates: A literature review. EPA-600/3-78-103. National Technical Information Service, Springfield, VA.

Pickering, Q.H. 1974. Chronic toxicity of nickel to the fathead minnow. J. Water Pollut. Control Fed. 46:760-765.

Pickering, Q.H. and C. Henderson. 1966. The acute toxicity of some heavy metals to different species of warmwater fishes. Air Water Pollut. Int. J. 10:453-463.

Portmann, J.E. 1968. Progress report on a programme of insecticide analysis and toxicity-testing in relation to the marine environment. Helgol. Wiss. Meeresunters. 17:247-256.

Portmann, J.E. 1972. Possible dangers of marine pollution as a result of mining operations for metal ores. In: Marine pollution and sea life. Ruivo, M. (Ed.). Fishing News (Books) Ltd., Surrey, England. pp. 343-346.

Pringle, B.H., D.E. Hissong, E.L. Katz and S.T. Mulawka. 1968. Trace metal accumulation by estuarine molluscs. J. Sanit. Eng. Div. Proc. Am. Soc. Civ. Eng. 94:455-475.

Pulich, W.M. 1980. Heavy metal accumulation by selected Halodule wrightii Asch. populations in the Corpus Christi Bay area. Contrib. Mar. Sci. 23:89-100.

Rai, L.C. and M. Raizada. 1985. Effect of nickel and silver ions on survival, growth, carbon fixation and nitrogenase activity in Nostoc muscorum: Regulation of toxicity by EDTA and calcium. J. Gen. Appl. Microbiol. 31:329-337.

Rai, L.C., J.P. Gaur and H.D. Kumar. 1981. Phycology and heavy-metal pollution. *Ecol. Rev. Camb. Philos. Soc.* 56:99-151.

Rehwoldt, R., G. Bida and B. Nerrie. 1971. Acute toxicity of copper, nickel and zinc ions to some Hudson River fish species. *Bull. Environ. Contam. Toxicol.* 6:445-448.

Rehwoldt, R., L.W. Menapace, B. Nerrie and D. Alessandrello. 1972. The effect of increased temperature upon the acute toxicity of some heavy metal ions. *Bull. Environ. Contam. Toxicol.* 8:91-96.

Rehwoldt, R., L. Lawrence, C. Shaw and E. Wirhowski. 1973. The acute toxicity of some heavy metal ions toward benthic organisms. *Bull. Environ. Contam. Toxicol.* 10:291-294.

Reynolds, B.H. 1979. Trace metals monitoring at two ocean disposal sites. EPA-600/3-79-037. National Technical Information Service, Springfield, VA.

Sato, C., J.L. Schnoor and D.B. McDonald. 1986. Characterization of effects of copper, cadmium and nickel on the growth of Nitrosomonas europaea. *Environ. Toxicol. Chem.* 5:403-416.

Saxena, O.P. and A. Parashari. 1983. Comparative study of the toxicity of six heavy metals to Channa punctatus. *J. Environ. Biol.* 4:91-94.

See, C.L., A.L. Buikema and J. Cairns. 1974. The effects of selected toxicants on survival of Dugesia tigrina (Turbellaria). Assoc. Southeastern Biologists Bull. 21:82.

See, C.L., A.L. Buikema and J. Cairns. 1975. The effects of sublethal concentrations of zinc and nickel on the photonegative response of Dugesia tigrina. Va. J. Sci. 26:60.

Seelye, J.G., R.J. Hesselberg and M.J. Mac. 1982. Accumulation by fish of contaminants released from dredged sediments. Environ. Sci. Technol. 16:459-464.

Shaw, T.L. and V.M. Brown. 1971. Heavy metals and the fertilization of rainbow trout eggs. Nature 230:251.

Shaw, W.H. and B. Grushkin. 1957. The toxicity of metal ions to aquatic organisms. Arch. Biochem. Biophys. 67:447-452.

Shcherban, E.P. 1977. Toxicity of some heavy metals for Daphnia magna Strauss, as a function of temperature. Hydrobiol. J. 13(4):75-80.

Sirover, M.A. and L.A. Loeb. 1976. Metal activation of DNA synthesis. Biochem. Biophys. Res. Commun. 7:812-817.

Skaar, H., B. Rystad and A. Jensen. 1974. The uptake of ^{63}Ni by the diatom Phaeodactylum tricornutum. Physiol. Plant. 32:353-358.

Smith-Sonneborn, J., R.A. Palizzi, E.A. McCann and G.L. Fisher. 1983.

Bioassay of genotoxic effects of environmental particles in a feeding ciliate. Environ. Health Perspect. 51:205-210.

Soeder, C.J. and G. Engelmann. 1984. Nickel requirement in Chlorella emersonii. Arch. Microbiol. 137:85-87.

Solbe, J.F. 1973. The relation between water quality and the status of fish populations in Willow Brook. Water Treat. Exam. 22:41-58.

Spencer, D.F. and R.W. Greene. 1981. Effects of nickel on seven species of freshwater algae. Environ. Pollut. (Series A) 25:241-247.

Spencer, D.F. and L.H. Nichols. 1983. Free nickel ion inhibits growth of two species of green algae. Environ. Pollut. (Series A) 31:97-104.

Sprague, J.B. 1985. Factors that modify toxicity. In: Fundamentals of aquatic toxicology - methods and applications. Rand, G.M. and S.R. Petrocelli (Eds.). Hemisphere Publishing Corporation, New York, NY. pp. 124-163.

Srivastava, D.K., S.S. Khanna, V.M. Srivastwa and R. Kumar. 1985. Electron diffraction of the metallic elements in the pineal organ of a freshwater fish, Mystus vittatus (Bloch). Experimentia 41:603-605.

Stephan, C.E., D.I. Mount, D.J. Hansen, J.H. Gentile, G.A. Chapman and W.A. Brungs. 1985. Guidelines for deriving numerical national water quality criteria for the protection of aquatic organisms and their uses. PB85-227049. National Technical Information Service, Springfield, VA.

Stokes, P. 1975. Uptake and accumulation of copper and nickel by metal-tolerant strains of Scenedesmus. Int. Ver. Theor. Angew. Limnol. Verh. 19:2128-2137.

Stokes, P.M., T.C. Hutchinson and K. Krauter. 1973. Heavy-metal tolerance in algae isolated from contaminated lakes near Sudbury, Ontario. Can. J. Bot. 51:2155-2168.

Stokes, P.M., R.C. Bailey and G.R. Groulx. 1985. Effects of acidification on metal availability to aquatic biota, with special reference to filamentous algae. Environ. Health Perspect. 63:79-87.

Stratton, G.W. and C.T. Corke. 1979a. The effect of nickel on the growth, photosynthesis, and nitrogenase activity of Anabaena inaequalis. Can. J. Microbiol. 25:1094-1099.

Stratton, G.W. and C.T. Corke. 1979b. The effect of mercuric, cadmium, and nickel ion combinations on a blue-green alga. Chemosphere 10:731-740.

Suloway, J.J., W.R. Roy, T.M. Skelly, D.R. Dickerson, R.M. Schuller and R.A. Griffin. 1983. Chemical and toxicological properties of coal fly ash. Illinois State Geological Survey Division, Champaign, IL.

Swaine, D.J. 1980. Nickel in coal and fly ash. In: Nickel in the environment. Nriagu, J.O. (Ed.). Wiley, New York, NY. pp. 67-92.

Tarzwell, C.M. and C. Henderson. 1960. Toxicity of less common metals to fishes. Ind. Wastes 5:12.

Taylor, G.J. and A.A. Crowder. 1983. Uptake and accumulation of copper, nickel and iron by Typha latifolia grown in solution culture. Can. J. Bot. 61:1825-1830.

Taylor, G.J. and A.A. Crowder. 1984. Copper and nickel tolerance in Typha latifolia clones from contaminated and uncontaminated environments. Can. J. Bot. 62:1304-1308.

Thompson, S.E., C.A. Burton, D.J. Quinn and Y.C. Ng. 1972. Concentration factors of chemical elements in edible aquatic organisms. UCRL-50564. Rev. 1. National Technical Information Service, Springfield, VA.

Timourian, H. and G. Watchmaker. 1972. Nickel uptake by sea urchin embryos and their subsequent development. J. Exp. Zool. 182:379-388.

Timourian, N. and G. Watchmaker. 1977. Assay of sperm motility to study the effects of metal ions. In: Biological implications of metals in the environment. ERDA Symposium Series 42.

Tong, S.S., W.D. Youngs, W.H. Gotenmann and D.J. Lisk. 1974. Trace metals in Lake Cayuga lake trout (Salvelinus namaycush) in relation to age. J. Fish. Res. Board Can. 31:238-239.

Trollope, D.R. and B. Evans. 1976. Concentrations of copper, iron, lead, nickel and zinc in freshwater algal blooms. Environ. Pollut. 11:109-116.

U.S. EPA. 1975. Preliminary investigation of effects on the environment of boron, indium, nickel, selenium, tin, vanadium and their compounds.

EPA-560/2-75-005C. National Technical Information Service, Springfield, VA.

U.S. EPA. 1976. Quality criteria for water. EPA-440/9-76-023. National Technical Information Service, Springfield, VA.

U.S. EPA. 1980. Ambient water quality criteria for nickel. EPA-440/5-80-060. National Technical Information Service, Springfield, VA.

U.S. EPA. 1983a. Methods for chemical analysis of water and wastes. EPA-600/4-79-020 (Revised March 1983). National Technical Information Service, Springfield, VA.

U.S. EPA. 1983b. Water quality standards regulation. Fed. Regist. 48:51400-51413. November 8.

U.S. EPA. 1983c. Water quality standards handbook. Office of Water Regulations and Standards, Washington, DC.

U.S. EPA. 1985a. Appendix B - Response to public comments on "Guidelines for deriving numerical national water quality criteria for the protection of aquatic organisms and their uses." Fed. Regist. 50:30793-30796. July 29.

U.S. EPA. 1985b. Technical support document for water-quality based toxics control. Office of Water, Washington, DC. September.

U.S. EPA. 1986. Chapter 1 - Stream design flow for steady-state modeling. In: Book VI - Design conditions. In: Technical guidance manual for performing waste load allocations. Office of Water, Washington, DC.

Uthe, J.F. and E.G. Bligh. 1971. Preliminary survey of heavy metal contamination of Canadian freshwater fish. J. Fish. Res. Board Can. 28:786-788.

Van Coillie, R. and A. Rousseau. 1974. Mineral composition in scales of Catostromus commersoni exposed to different media: Analysis by micro-electron technique. J. Fish. Res. Board Can. 31:63-66.

Van Hoof, F. and J.P. Nauwelaers. 1984. Distribution of nickel in the roach (Rutilus rutilus L.) after exposure to lethal and sublethal concentrations. Chemosphere 13:1053-1058.

Verma, S.R., M. Jain and R.C. Dalela. 1981. In vivo removal of a few heavy metals in certain tissues of the fish Notopterus notopterus. Environ. Res. 26:328-334.

Vymazal, J. 1984. Short-term uptake of heavy metals by periphyton algae. Hydrobiologia 119:171-179.

Wachs, V.B. 1982. Concentration of heavy metals in fishes from the river Danube. Z. Wasser Abwasser Forsch 15:43-49.

Wang, H.K. and J.M. Wood. 1984. Bioaccumulation of nickel by algae. Environ. Sci. Technol. 18:106-109.

Wang, W. 1986. Toxicity tests of aquatic pollutants by using common duckweed. Environ. Pollut. (Series B) 11:1-14.

Wang, Z., C.P. Bianchi and S.R. Narayan. 1984. Nickel inhibition of calcium release from subsarcolemmal calcium stores of molluscan smooth muscle. J. Pharmacol. Exp. Ther. 229:696-701.

Warnick, S.L. and H.L. Bell. 1969. The acute toxicity of some heavy metals to different species of aquatic insects. J. Water Pollut. Control Fed. 41:280-284.

Waterman, A.J. 1937. Effect of salts of heavy metals on the development of the sea urchin, *Arbacia punctulata*. Biol. Bull. (Woods Hole) 73:401-420.

Watling, H.R. 1983. Comparative study of the effects of metals on the settlement of *Crassostrea gigas*. Bull. Environ. Contam. Toxicol. 31:344-351.

Watras, C.J., J. MacFarlane and F.M. Morel. 1985. Nickel accumulation by Scenedesmus and Daphnia: Food-chain transport and geochemical implications. Can. J. Fish. Aquat. Sci. 42:724-730.

Weber, W.J., Jr. and W. Stumm. 1963. Mechanisms of hydrogen ion buffering in natural waters. J. Am. Water Works Assoc. 55:1553-1578.

Wehr, J.D. and B.A. Whitton. 1983. Accumulation of heavy metals by aquatic mosses. 2: *Rhynchostegium riparioides*. Hydrobiologia 100:261-284.

Weinstein, N.L. and P.D. Anderson. 1978. Lethal and sublethal toxicities of copper-nickel mixtures to the zebrafish *Brachydanio rerio*. In: Proceedings of the fourth annual aquatic toxicity workshop. Davis, J.D., G.L. Greer and I.K. Birtwell (Eds.). Fisheries and Marine Service Technical Report No. 818. Department of Fisheries and Oceans, Ottawa, Ontario, Canada. p. 154.

Whitley, L.S. and R.A. Sikora. 1970. The effect of three common pollutants on the respiration rate of tubificid worms. J. Water Pollut. Control Fed. 42:R57-R66.

Whitton, B.A. and F.H.A. Shehata. 1982. Influence of cobalt, nickel, copper and cadmium on the blue-green alga Anacystis nidulans. Environ. Pollut. (Series A) 27:275-281.

Willford, W.A. 1966. Toxicity of 22 therapeutic compounds to six fishes. Investigations in Fish Control 18. U.S. Fish and Wildlife Service, Washington, DC.

Wilson, J.G. 1983. The uptake and accumulation of Ni by Cerastoderma edule and its effect on mortality, body condition and respiration rate. Mar. Environ. Res. 8:129-148.

Wilson, W.B. and L.R. Freeberg. 1980. Toxicity of metals to marine phytoplankton cultures. EPA-600/3-80-025. National Technical Information Service, Springfield, VA.

Windom, H.L., K. Tenore and D.L. Rice. 1982. Metal accumulation of the polychaete Capitella capitata: Influences of metal content and nutritional quality of detritus. Can. J. Fish. Aquat. Sci. 39:191-196.

Wong, P.T., Y.K. Chau and P.L. Luxon. 1978. Toxicity of a mixture of metals on freshwater algae. J. Fish. Res. Board Can. 35:479-481.

Wong, P.T.S., Y.K. Chau and D. Patel. 1982. Physiological and biochemical responses of several freshwater algae to a mixture of metals. Chemosphere 11:367-376.

Wong, S.L. and J.L. Beaver. 1980. Algal bioassays to determine toxicity of metals mixtures. *Hydrobiologia* 74:199-208.

Wren, C.D., H.R. MacCrimmon and B.R. Loescher. 1983. Examination of bioaccumulation and biomagnification of metals in a precambrian shield lake. *Water Air Soil Pollut.* 19:277-291.

Yan, N.D., G.E. Miller, I. Wile and G.G. Hitchin. 1985. Richness of aquatic macrophyte floras of soft water lakes of differing pH and trace metal content in Ontario, Canada. *Aquat. Bot.* 23:27-40.

Young, D.R. 1982. A comparative study of trace metal contamination of the Southern California and New York Bights. In: *Ecological stress and the New York Bight: Science and management*. Mayer, G.F. (Ed.). Estuarine Research Federation, Columbia, SC. pp. 249-262.

Zarogian, G.E. and M. Johnson. 1984. Nickel uptake and loss in the bivalves Crassostrea virginica and Mytilus edulis. *Arch. Environ. Contam. Toxicol.* 13:411-418.

Zarogian, G.E., J.H. Gentile, J.F. Heltsche, U. Johnson and A.M. Ivanovici. 1982. Application of adenine nucleotide measurements for the evaluation of stress in Mytilus edulis and Crassostrea virginica. *Comp. Biochem. Physiol.* 71B:643-649.